

Invertible Dirac operators and handle attachments

Nadine Große

Universität Leipzig

(joint work with Mattias Dahl)

Rauischholzhausen, 03.07.2012

Motivation

- ▶ Not every closed manifold admits a metric of positive scalar curvature.
- ▶ In contrast, on every closed manifold the space of metric with negative scalar curvature is nonempty and contractible.
- ▶ Topological obstruction for psc-metrics:
(M, g) closed spin, Dirac operator D^g

Lichnerowicz formula

$$(D^g)^2 = \Delta_g + \frac{\text{scal}_g}{4}$$

$\text{scal}_g > 0 \Rightarrow D^g$ is invertible

Motivation

- ▶ Not every closed manifold admits a metric of positive scalar curvature.
- ▶ Topological obstruction for psc-metrics:
(M, g) closed spin, Dirac operator D^g

Lichnerowicz formula

$$(D^g)^2 = \Delta_g + \frac{\text{scal}_g}{4}$$

$\text{scal}_g > 0 \Rightarrow D^g$ is invertible

- ▶ $\text{Metr}(M)^{\text{psc}} \subset \text{Metr}(M)^{\text{inv}} \subset \text{Metr}(M)$

Obstruction for psc metrics

From index theory

$$\dim \ker D^g \geq \begin{cases} |\hat{A}(M)| & \text{if } n \equiv 0 \pmod{4} \\ 1 & \text{if } n \equiv 1 \pmod{8}, \alpha(M) \neq 0 \\ 2 & \text{if } n \equiv 2 \pmod{8}, \alpha(M) \neq 0 \\ 0 & \text{otherwise} \end{cases}$$



where \hat{A} and α are determined only by the topology of the underlying manifold.

E.g. if $\hat{A}(M^4) \neq 0$, $\text{Metr}(M^4)^{\text{psc}} \subset \text{Metr}(M^4)^{\text{inv}} = \emptyset$.

$\text{Metr}(M)^{\text{psc}} \subset \text{Metr}(M)^{\text{inv}} \subset \text{Metr}(M)$

Construction of manifolds admitting psc-metrics

- Review Surgery




- ▶ embedding $f : S^k \times B^{n-k} \rightarrow M$
 $S := f(S^k \times \{0\})$ - surgery sphere
- ▶ $\partial(M \setminus f(S^k \times B^{n-k})) \cong S^{k-1} \times S^{n-k-1}$
- ▶ $M' = (M \setminus f(S^k \times B^{n-k})) \sqcup_{\sim} B^{k+1} \times S^{n-k-1}$

M' is obtained from M by a surgery of $\dim k$ / $\text{codim } n - k$.

Construction of manifolds admitting psc-metrics

- Review Surgery




- ▶ View the cylinder $W := M \times [0, 1]$ as a bordism from M to itself
- ▶ Attach $B^{k+1} \times B^{n-k-1}$ to $M \times \{1\}$
- ▶ W' is a bordism from M to M' - the trace of the surgery.

W' is obtained from W by attaching a $(k + 1)$ -handle.

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.




The torus is obtained as:

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.




The torus is obtained as:

$$B^2 +$$

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.



$$B^1 \times B^1$$

$$B^2$$


The torus is obtained as:

$$B^2 + \text{a 1-handle}$$

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.



$$B^1 \times B^1$$
$$B^2$$



The torus is obtained as:

$$B^2 + \text{a 1-handle}$$

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.





The torus is obtained as:

$$B^2 + \text{a 1-handle}$$

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.





The torus is obtained as:

$$B^2 + \text{a 1-handle} + \text{a 1-handle}$$

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.





The torus is obtained as:

$$B^2 + \text{a 1-handle} + \text{a 1-handle}$$

Construction of manifolds admitting psc-metrics

- Review Surgery

Each closed manifold has a handle decomposition.





The torus is obtained as:

$$B^2 + \text{a 1-handle} + \text{a 1-handle} + B^2 = T^2$$

Construction of manifolds admitting psc-metrics



Theorem (Gromov, Lawson / Schoen, Yau; '80)

Let (M, g) be a closed Riemannian manifold with $g \in \text{Metr}(M)^{\text{psc}}$.
Let M' be obtained from M by a surgery of codimension ≥ 3 .
Then, M' admits a psc-metric g' .



g' can be chosen such that it coincides with g outside a small neighbourhood around the surgery sphere.

Construction of manifolds admitting psc-metrics





Intuition

- ▶ psc is a local property
- ▶ $\text{codim } n - k \geq 3 = \text{gluing in } B^{k+1} \times S^{n-k-1} \geq 2$
- ▶ standard product structure on $B^{k+1} \times S^{n-k-1} \geq 2$ has psc



Psc-metrics and handle attachments

Theorem (Carr '88 / Gajer '87)

Let (M^{n+1}, g) be a compact Riemannian manifold with closed boundary ∂M , $g \in \text{Metr}(M)^{\text{psc}}$ and g having product structure near ∂M . Let M' be obtained from M by adding a $(k+1)$ -handle of codimension $n-k \geq 3$. Then, M' admits a psc-metric g' that is again product near the (new) boundary.




Psc-metrics and handle attachments



Intuition

- ▶ On the boundary: surgery of codim $n - k \geq 3$


Psc-metrics and handle attachments



Intuition

- ▶ On the boundary: surgery of codim $n - k \geq 3$
- ▶ On the double: surgery of codim $n - k \geq 3$

Psc-metrics and handle attachments



Intuition

- ▶ On the boundary: surgery of codim $n - k \geq 3$
- ▶ On the double: surgery of codim $n - k \geq 3$

Implication

- ▶ $\text{Metr}^{\text{psc}}(S^{4k-1})$ has infinitely many components ($k \geq 2$)
($\text{Metr}^{\text{psc}}(S^3)$ is connected (Marques, 2011))

Metr^{inv}(M)



What can be done for metrics with invertible Dirac operators?

Metr^{inv}(M)

What can be done for metrics with invertible Dirac operators?



From now on: Let all manifolds be spin.

Surgery for $\text{Met}^{\text{inv}}(M)$





- After the surgery the manifold should still be spin!

Surgery for $\text{Met}^{\text{inv}}(M)$





- ▶ After the surgery the manifold should still be spin!
 - ▶ $S^k \times B^{n-k}$ - induces spin structure on $S^k \times S^{n-k-1}$
 - ▶ glue in $B^{k+1} \times S^{n-k-1}$
 - Its boundary should carry same spin structure.
 - ▶ For $k > 1$, the spin structure on S^k is unique and bounds the disk. - No Problem here.
 - ▶ For $k = 1$, two spin structures on S^1 - we only allow the one that bounds the disk.

Surgery for $\text{Met}^{\text{inv}}(M)$



- ▶ $f : S^k \times B^{n-k} \rightarrow M$ spin-preserving embedding.



Surgery for $\text{Met}^{\text{inv}}(M)$



Intuition

- ▶ Invertible Dirac operator is a **global** condition.
- ▶ $\text{codim } n - k \geq 3 = \text{gluing in } B^{k+1} \times S^{n-k-1} \geq 2$
- ▶ standard product structure on $\mathbb{R}^{k+1} \times S^{n-k-1} \geq 2$ has invertible Dirac operator



Surgery for $\text{Met}^{\text{inv}}(M)$



Intuition

- ▶ Invertible Dirac operator is a **global** condition.
- ▶ $\text{codim } n - k \geq 2 = \text{gluing in } B^{k+1} \times S^{n-k-1} \geq 1$
- ▶ standard product structure on $\mathbb{R}^{k+1} \times S^{n-k-1} \geq 1$ has invertible Dirac operator ('When taking the right S^1 ')

Surgery for $\text{Met}^{\text{inv}}(M)$



Intuition

- ▶ Invertible Dirac operator is a **global** condition.
- ▶ $\text{codim } n - k \geq 2$ = gluing in $B^{k+1} \times S^{n-k-1} \geq 1$
- ▶ standard product structure on $\mathbb{R}^{k+1} \times S^{n-k-1} \geq 1$ has invertible Dirac operator ('When taking the right S^1 ')
- ▶ 'If the inserted cylinder is large enough, invertibility survives.'

Construction for manifolds admitting inv-metrics


Theorem (Ammann, Dahl, Humbert; 2009)

Let (M^n, g) be a closed Riemannian spin manifold with $g \in \text{Metr}(M)^{\text{inv}}$. Let M' be obtained from M by a surgery of codimension ≥ 2 . Then, M' admits an inv-metric g' . Moreover, g' can be chosen such that it coincides with g outside a small neighbourhood around the surgery sphere.

Consequences (Ammann, Dahl, Humbert; 2009)


For a generic metric g , $\dim \ker D^g$ is no larger than forced by the index theorem.

Inv-metrics on manifolds with boundary




When do we call D^g invertible?

Inv-metrics on manifolds with boundary




$g \in \text{Metr}(M)^{\text{inv}}$ iff D^{g_∞} is invertible as operator on $L^2(M_\infty, S)$

Inv-metrics on manifolds with boundary



$$(M_1, g_1), (\partial M_1, \partial g_1) = (N^+, h),$$

$$g_1 \in \text{Metr}(M_1)^{\text{inv}}$$



$$(M_2, g_2),$$



$$(\partial M_2, \partial g_2) = (N^-, h), g_2 \in \text{Metr}(M_2)^{\text{inv}}$$

If M_1 and M_2 are glued together using a **large enough cylinder** $(N \times [-R, R], h + dt^2)$, the resulting metric has again invertible Dirac operator.

Inv-metrics and handle attachments

Theorem (Dahl, G. 2012)

Let (M^{n+1}, g) be a compact Riemannian spin manifold with closed boundary ∂M , $g \in \text{Metr}(M)^{\text{inv}}$ and g having product structure near ∂M . Let M' be obtained from M by adding a $(k+1)$ -handle of codimension $n-k \geq 2$. Then, M' admits an inv-metric g' that is again product near the (new) boundary.



Inv-metrics and handle attachments

Theorem (Dahl, G. 2012)



Let (M^{n+1}, g) be a compact Riemannian spin manifold with closed boundary ∂M , $g \in \text{Metr}(M)^{\text{inv}}$ and g having product structure near ∂M . Let M' be obtained from M by adding a $(k+1)$ -handle of codimension $n - k \geq 2$. Then, M' admits an inv-metric g' that is again product near the (new) boundary.

Implication



- ▶ $\text{Metr}(S^{4k-1})^{\text{inv}}$ has infinitely many components for all $k \geq 1$

Strategy and Methods

- 'Topological strategy' - Decompose the handle attachment



surgery of codim $n - k$






'half' surgery of codim $n - k + 1$
glue in ' $\frac{1}{2}B^{k+1} \times S^{n-k}$ '

Metric strategy

- Approximation by 'double' product metrics near the surgery sphere

$$(U_\delta(S \times [-\epsilon, \infty)),$$

$$g_S + \xi_{\mathbb{R}^{n-k}} + dt^2)$$




$$(\partial M \times [-\epsilon, \infty), \\ , \partial g + dt^2)$$

$$\leftarrow \partial M \times \{0\}$$

If δ small enough, still $g_\delta \in \text{Metr}(M)^{\text{inv}}$.
(C^1 -continuity of the spectrum')

Metric strategy


► First surgery



- 'Parameter for tuning': ρ - 'diameter of $\{B\}$ '
- For ρ small enough, $g'_\rho \in \text{Metr}(M')^{\text{inv}}$ - proof by contradiction

Metric strategy


► First surgery



- For ρ small enough, $g_\rho \in \text{Metr}(M')^{\text{inv}}$ - proof by contradiction
 - $\rho_i \rightarrow 0$, $g_{\rho_i} \notin \text{Metr}(M')^{\text{inv}}$
 - $\leadsto g_{\rho_i}$ has a harmonic spinor: $D^{g_{\rho_i}} \varphi_i = 0$, $\|\varphi_i\|_{L^2(M', g_{\rho_i})} = 1$
 - (regularity) $\leadsto \varphi_i \rightarrow \varphi$ in $C^1_{\text{loc}}(M \setminus (S \times [-\epsilon, \infty)))$
 - (removal of singularities) $\leadsto D^g \varphi = 0$ on M , $\|\varphi\|_{L^2(M, g)} \leq 1$
- a priori estimates on the L^2 -norm of φ_i on $\{A\}$ vs $\{C\}$.


Metric strategy

- ▶ Again approximating by 'double' product metrics



Metric strategy

- ▶ Second surgery



One Application



Theorem (Dahl, G.; 2012)

Let M be a closed 3-dimensional Riemannian spin manifold and $g \in \text{Metr}(M)^{\text{inv}}$. Then there are metrics $g^i \in \text{Metr}(M)^{\text{inv}}$, $i \in \mathbb{Z}$, such that g^i is bordant to g but g^i is not concordant to g^j for $i \neq j$.

In particular, $\text{Metr}(M)^{\text{inv}}$ has infinitely many connected components.

Notations


$g_0, g_1 \in \text{Metr}(M)^{\text{inv}}$ are *isotopic* if \exists smooth family $g_t \in \text{Metr}(M)^{\text{inv}}$ with $g_t = g_0$ for $t \leq 0$, $g_t = g_1$ for $t \geq 1$.



(M, g_0) (M, g_1)

$g_0, g_1 \in \text{Metr}(M)^{\text{inv}}$ are *concordant* if $\exists \tilde{g} \in \text{Metr}(M \times [0, 1])^{\text{inv}}$ with $\tilde{g}|_{M \times \{i\}} = g_i$.

$g_i \in \text{Metr}(M_i)^{\text{inv}}$ ($i = 0, 1$) are *bordant* if $\exists (W, \tilde{g})$ with $\partial W = M_0 \sqcup (M_1)^-$, $\tilde{g} \in \text{Metr}(W)^{\text{inv}}$, $\tilde{g}|_{M_i} = g_i$.



(M_0, g_0) (M_1, g_1)

An application

Theorem (Dahl, G.; 2012)

Let M be a closed 3-dimensional Riemannian spin manifold and $g \in \text{Metr}(M)^{\text{inv}}$. Then there are metrics $g^i \in \text{Metr}(M)^{\text{inv}}$, $i \in \mathbb{Z}$, such that g^i is bordant to g but g^i is not concordant to g^j for $i \neq j$.

In particular, $\text{Metr}(M)^{\text{inv}}$ has infinitely many connected components.

Lemma

There exist 4-manifolds (Y^i, \tilde{h}^i) ($i \in \mathbb{Z}$) with $\tilde{h}^i \in \text{Metr}(Y^i)^{\text{inv}}$, $\partial Y^i = S^3$ such that $\alpha(Y^i \cup_{S^3} (Y^j)^-) = c(i - j)$ for a constant $c \neq 0$.

An application

Lemma

There exist 4-manifolds (Y^i, \tilde{h}^i) ($i \in \mathbb{Z}$) with $\tilde{h}^i \in \text{Metr}(Y^i)^{\text{inv}}$, $\partial Y^i = S^3$ such that $\alpha(Y^i \cup_{S^3} (Y^j)^-) = c(i-j)$ for a constant $c \neq 0$.

Construction:


- ▶ $Y^0 - B^4$ with a 'torpedo metric' $\tilde{h}^0 \in \text{Metr}(B^4)^{\text{psc}}$ and $\tilde{h}^0|_{S^3} = \text{standard metric}$
- ▶ $Y^i = \underbrace{(K3 \# K3 \# \dots \# K3)}_{i \text{ times}} \setminus B^4 = Y^0 + \text{several 2-handles}$
- ▶ $\alpha(Y^i \cup_{S^3} (Y^j)^-) = \alpha(\#_{(i-j)} K3) = (i-j)\alpha(K3) \neq 0$ for $i \neq j$

$$h^i := \tilde{h}^i|_{S^3}$$

Constructions of g^i

Start with $(M^3, g \in \text{Metr}(M)^{\text{inv}})$ - construct g^i bordant to g .

(M, g)




(M, g^i)

(S^3, h^i)


attachment of a 1-handle

Constructions of g^i




(M, g) and (M, g^i) are bordant.
Bordism $(W^i, \tilde{g}^i) \in \text{Metr}(W^i)^{\text{inv}}$

Constructions of g^i



Constructions of g^i



Closed manifold (W, \tilde{g}) with $\tilde{g} \in \text{Metr}(W)^{\text{inv}}$ and $\alpha(W) = c(i - j)$.

One Application

Theorem (Dahl, G.; 2012)

Let M be a closed 3-dimensional Riemannian spin manifold and $g \in \text{Metr}(M)^{\text{inv}}$. Then there are metrics $g^i \in \text{Metr}(M)^{\text{inv}}$, $i \in \mathbb{Z}$, such that g^i is bordant to g but g^i is not concordant to g^j for $i \neq j$.

In particular, $\text{Metr}(M)^{\text{inv}}$ has infinitely many connected components.

Thank you for your attention.