

$$i\mathbb{L}_{\infty}^0 \mathbb{C} \subset i\mathbb{L}_{\infty}^1 \mathbb{C} \subset i\mathbb{L}_{\infty}^2 \mathbb{C}$$

$$\gamma \star \acute{\gamma} = \int_{dx/(2\pi)^d}^{\mathbb{L}} ix^{-} ix^{\acute{\gamma}}$$

schnell fallend $i\mathbb{L}_{\infty}^1 \mathbb{C} = \frac{\gamma \in i\mathbb{L}_{\infty}^1 \mathbb{C}}{p \cdot \partial_q \gamma \text{ bes } \wedge p:q \in i\mathbb{L}_{\infty}^1 \mathbb{C}}$

$$i\mathbb{R}^d_{\infty}^0 \mathbb{C} \subset i\mathbb{R}^d_{\infty}^1 \mathbb{C} \subset i\mathbb{R}^d_{\infty}^2 \mathbb{C}$$

$$\gamma \star \acute{\gamma} = \int_{dx/(2\pi)^d}^{\mathbb{R}^d} i\mathbb{L}^{-} i\mathbb{L}^{\acute{\gamma}}$$

schnell fallend $i\mathbb{R}^d_{\infty}^1 \mathbb{C} = \frac{\gamma \in i\mathbb{R}^d_{\infty}^1 \mathbb{C}}{p \cdot \partial_q \gamma \text{ bes } \wedge p:q \in i\mathbb{R}^d_{\infty}^1 \mathbb{C}}$