

Jacobs University Bremen
School of Engineering and Science
Götz Pfander

Fall Term 2007,
Spring Term 2008

Analysis I,II

Version (December 6, 2007)

Contents

1	Numbers	3
1.1	Sets, relations and functions	3
1.2	Groups, fields, the integers and the rational numbers	6
1.3	Real numbers	9
1.4	Complex numbers	13
2	Convergence of sequences in metric spaces and numeric series	14
2.1	Sequences in metric spaces	14
2.2	The extended real number system, \limsup and \liminf	17
2.3	Cauchy sequences and complete metric spaces	19
2.4	Real and complex series	20
3	Topology and continuity	25
3.1	Continuous functions	25
3.2	Topological spaces	27
3.3	Compactness	30
3.4	Connectedness	34
3.5	Sequences of functions, uniform convergence	36
4	Differentiation	37
4.1	Central results	37
4.2	Taylor series	40
4.3	The exponential function and friends	43
4.4	Fixed point theorems and approximative methods	45

Preface

This script contains all the theorems and definitions, but only a few examples covered in Analysis I, II in the academic year 2007/2008.

Most proofs have been omitted from this script. With the exception of two or three theorems, all statements have been proven in either the script, in class, or in the homeworks.

1. NUMBERS

1.1. Sets, relations and functions

DEFINITION 1.1. The *cartesian product* $X_1 \times X_2 \times \dots \times X_n$ of the n sets X_1, X_2, \dots, X_n is the set of all (ordered) *n-tuples* (x_1, x_2, \dots, x_n) with $x_1 \in X_1, x_2 \in X_2, \dots, x_n \in X_n$. That is,

$$X_1 \times X_2 \times \dots \times X_n := \{(x_1, x_2, \dots, x_n) : x_1 \in X_1, x_2 \in X_2, \dots, x_n \in X_n\}.$$

Note that $A \times \emptyset = \emptyset \times A = \emptyset$, and $A \times B = B \times A$ if and only if $A = B$ or $A = \emptyset$ or $B = \emptyset$.

EXAMPLES 1.2.

- i. $\{1, 2, 3\} \times \{7, 12\} = \{(1, 7), (2, 7), (3, 7), (1, 12), (2, 12), (3, 12)\}$
- ii. $\{7, 12\} \times \{1, 2, 3\} = \{(7, 1), (7, 2), (7, 3), (12, 1), (12, 2), (12, 3)\}$

DEFINITION 1.3. Any subset R of the cartesian product $X \times Y$ of two sets X and Y , that is, $R \subset X \times Y$, is called *relation* between X and Y . If $X = Y$ we say that $R \subset X \times X$ is a relation on X .

$\mathcal{D}(R) = \mathcal{D}_R = \{x \in X : \text{there exists } y \in Y \text{ with } (x, y) \in R\}$ is called *domain of R*, and
 $\mathcal{R}(R) = \mathcal{R}_R = \{y \in Y : \text{there exists } x \in X \text{ with } (x, y) \in R\}$ is called *range of R*.

DEFINITION 1.4. Let X and Y be sets. A *function* (or *mapping*) $f : X \longrightarrow Y$ is a rule that associates to **every** element in $x \in X$ an element $f(x) \in Y$. X is called *domain* of f and is denoted by \mathcal{D}_f .

For $A \subseteq X$ and $B \subseteq Y$ we set

$$f(A) = \{y \in Y : \text{there exists } x \in A \text{ with } f(x) = y\}$$

and

$$f^{-1}(B) = \{x \in X : \text{there exists } y \in B \text{ with } f(x) = y\}.$$

The *range* of f is given by $\mathcal{R}_f = f(X)$. The *graph* of f is the relation $\Gamma_f = \{(x, y) \in X \times Y : f(x) = y\}$ between X and Y .

The function f is *injective* (*one-to-one*) if $f(x) = f(\tilde{x})$ implies $x = \tilde{x}$, and f is *surjective* (*onto*) if $\mathcal{R}_f = Y$. If f is surjective and injective, we call f *bijective*.

REMARK 1.5. Note that the distinction between a function and its graph is done for psychological reasons only. A strictly axiomatic introduction of analysis is based on set theory and functions are simply defined as certain subsets of $X \times Y$.

PROPOSITION 1.6. A relation $\Gamma \subset X \times Y$ is the graph of a function $f : \mathcal{D}_\Gamma \longrightarrow Y$, if and only if $(x, y), (x, \tilde{y}) \in \Gamma$ implies $y = \tilde{y}$ for all $x \in X$ and $y, \tilde{y} \in Y$. In this case we have $\mathcal{R}_f = \mathcal{R}_{\Gamma_f}$ and $\mathcal{D}_f = \mathcal{D}_{\Gamma_f}$.

THEOREM 1.7. Given a function $f : X \longrightarrow Y$ and sets $A_i \subset X$, $i \in \mathbb{N}$, and $B_i \subset Y$, $i \in \mathbb{N}$, we have

- i. $A_1 \subseteq A_2$ implies $f(A_1) \subseteq f(A_2)$
- ii. $B_1 \subseteq B_2$ implies $f^{-1}(B_1) \subseteq f^{-1}(B_2)$
- iii. $A_1 \subseteq f^{-1}(f(A_1))$ and $B_1 \supseteq f(f^{-1}(B_1))$
- iv. $f(\bigcup_{i=1}^{\infty} A_i) = \bigcup_{i=1}^{\infty} f(A_i)$ and $f(\bigcap_{i=1}^{\infty} A_i) \subseteq \bigcap_{i=1}^{\infty} f(A_i)$

If f is injective we have in addition $A_1 = f^{-1}(f(A_1))$ and $f(\bigcap_{i=1}^{\infty} A_i) = \bigcap_{i=1}^{\infty} f(A_i)$ and if f is surjective $B_1 = f(f^{-1}(B_1))$.

DEFINITION 1.8. A relation R on X is called

- i. *reflexive* if for all $x \in X$ we have $(x, x) \in R$,
- ii. *transitive* if $(x, \tilde{x}) \in R$ and $(\tilde{x}, \tilde{\tilde{x}}) \in R$ implies $(x, \tilde{\tilde{x}}) \in R$,
- iii. *symmetric* if $(x, \tilde{x}) \in R$ implies $(\tilde{x}, x) \in R$, and
- iv. *antisymmetric* if $(x, \tilde{x}) \in R$ and $(\tilde{x}, x) \in R$ implies $x = \tilde{x}$.

DEFINITION 1.9. A reflexive, symmetric, and transitive relation R on X is called *equivalence relation*. If R is an equivalence relation we shall write $x \sim \tilde{x}$ if $(x, \tilde{x}) \in R$ and call x and \tilde{x} *equivalent* with respect to R .

$[x] = \{\tilde{x} \in X : (x, \tilde{x}) \in R\}$ is called *equivalence class* of x , and any $\tilde{x} \in [x]$ is called *representative* of $[x]$.

The concept of a partition of a set helps to understand equivalence classes and their equivalence relations.

DEFINITION 1.10. A family of sets $\{M_i : i \in I\}$ is a partition of the set $M \neq \emptyset$, if

- i. $\emptyset \neq M_i \subset M$ for $i \in I$,
- ii. $i \neq j$ implies $M_i \cap M_j = \emptyset$ for $i, j \in I$, and
- iii. $\bigcup_{i \in I} M_i = M$.

THEOREM 1.11. For a set $M \neq \emptyset$ we have:

- i. The distinct equivalence classes of an equivalence relation on M form a partition on M .
- ii. A partition $\{M_i : i \in I\}$ on M induces an equivalence relation on M via

$$a \sim b \quad \text{if and only if} \quad a, b \in M_{i_0} \text{ for some } i_0 \in I.$$

EXAMPLE 1.12. Fix $n \in \mathbb{N}$ and set $X = \mathbb{Z}$. The relation

$$R_{\mathbb{Z}_n} = \{(k, m) \in \mathbb{Z} \times \mathbb{Z} : k - m = l \cdot n \text{ for some } l \in \mathbb{Z}\}$$

is an equivalence relation. The set of equivalence classes is the group \mathbb{Z}_n of n elements with addition given by

$$[n] + [m] = [n + m].$$

To see this, you would have to check whether addition is well defined and you need to check all group properties (which are discussed in detail below.)

1.2. Groups, fields, the integers and the rational numbers

DEFINITION 1.13. A *group* is a set G , together with a binary law of composition $\mu : G \times G \longrightarrow G$ which satisfies the axioms G1, G2, and G3 given below. We shall write $xy := \mu(x, y)$.

(G1) *Associativity*: $(xy)z = x(yz)$ for all $x, y, z \in G$.

(G2) *Identity*: There exists an element $e \in G$ called *identity* such that $xe = ex = x$ for all $x \in G$.

(G3) *Inverses* : To each element $x \in G$ exists an element $y \in G$ called *inverse* of x with $xy = yx = e$. The inverse to x is denoted by x^{-1} .

A group is called *abelian* if μ is commutative, that is, if we have

(C) $xy = yx$ for all $x, y \in G$.

EXAMPLES 1.14.

i. Let $X = \mathbb{N} \times \mathbb{N}$ and define

$$R_{\mathbb{Z}} = \{((n, m), (\tilde{n}, \tilde{m})) \in (\mathbb{N} \times \mathbb{N}) \times (\mathbb{N} \times \mathbb{N}) : n + \tilde{m} = \tilde{n} + m\}.$$

$R_{\mathbb{Z}}$ is an equivalence relation. The set of equivalence classes $\mathbb{Z} := \{[(n, m)]\}$ equipped with

- $[(n, m)] +_{\mathbb{Z}} [(\tilde{n}, \tilde{m})] = [(n + \tilde{n}, m + \tilde{m})]$
- $[(n, m)] \cdot_{\mathbb{Z}} [(\tilde{n}, \tilde{m})] = [(n \cdot \tilde{n} + m \cdot \tilde{m}, n \cdot \tilde{m} + m \cdot \tilde{n})]$
- $-[(n, m)] = [(m, n)]$

is a ring¹, called the *ring of integers*. We can embed (map injectively) the naturals into this ring of equivalence classes via

$$i : \mathbb{N} \longrightarrow \mathbb{Z}, \quad n \mapsto n^* := [(n + 1, 1)].$$

This mapping is nice, since it respects addition and multiplication on \mathbb{N} , that is,

$$i(n + \tilde{n}) = i(n) +_{\mathbb{Z}} i(\tilde{n}), \text{ and } i(n \cdot \tilde{n}) = i(n) \cdot_{\mathbb{Z}} i(\tilde{n})$$

Hence, using an appropriate equivalence relation on $\mathbb{N} \times \mathbb{N}$, we have created a ring of equivalence classes which can be identified with the set of integers.² In the following, we will not make a distinction between a natural number n and its integer counterpart n^* . We shall use the common short hand notation $z = n - m = [(n, m)] \in \mathbb{Z}$. Note that $[(7, 3)] = [(10, 6)]$, since $7 + 6 = 3 + 10$, that is, $7 - 3 = 10 - 6$

¹Since we shall not use any rings in this course, we omit a definition of *rings*. Please consult a textbook.

²We only assume *a-priori* knowledge of the naturals. Similar to the attitude of Leopold Kronecker, 1823-1891, who supposedly said “God made the integers; all else is the work of man”.

ii. Let $X = \mathbb{Z} \times \mathbb{N}$ and define

$$R_{\mathbb{Q}} = \{((z, m), (\tilde{z}, \tilde{m})) \in (\mathbb{Z} \times \mathbb{N}) \times (\mathbb{Z} \times \mathbb{N}) : z \cdot \tilde{m} = \tilde{z} \cdot m\}.$$

$R_{\mathbb{Q}}$ is an equivalence relation. The set of equivalence classes $\{[(z, m)]\}$ equipped with

- $[(z, m)] +_{\mathbb{Q}} [(\tilde{z}, \tilde{m})] = [(z \cdot_{\mathbb{Z}} \tilde{m} + \tilde{z} \cdot_{\mathbb{Z}} m, m \cdot_{\mathbb{Z}} \tilde{m})]$
- $[(z, m)] \cdot_{\mathbb{Q}} [(\tilde{z}, \tilde{m})] = [(z \cdot_{\mathbb{Z}} \tilde{z}, m \cdot_{\mathbb{Z}} \tilde{m})]$

is a field³, called the field of *rational numbers*. Again, we can embed the integers in a natural way by setting

$$i : \mathbb{Z} \longrightarrow \mathbb{Q}, \quad z \mapsto z^* := [(z, 1)].$$

This embedding respects multiplication and addition, hence, we consider \mathbb{Z} as a subring of the ring (field) of equivalence classes we just defined. The field we defined is the field of rational numbers. From now on, we shall use them the way we are used to. Certainly, we shall write $r = \frac{z}{m} = [(z, m)] \in \mathbb{Q}$.

Starting from the natural numbers we have created the integers, from those we have created the rationals. Since the embeddings are canonical, we shall ignore its formalism and simply take

$$\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q}.$$

DEFINITION 1.15. A *field* is a set F on which two binary laws of composition, *addition* $'+'$ and *multiplication* $'\cdot'$ are defined with

(F1) $(F, +)$ is an abelian group. We shall denote the identity of $(F, +)$ as 0.

(F2) $(F \setminus \{0\}, \cdot)$ is an abelian group. The identity of $(F \setminus \{0\}, \cdot)$ is denoted by 1.

(F3) The *distributive law* holds, that is, $(x + y) \cdot z = xz + yz$ for all $x, y, z \in F$.

DEFINITION 1.16. A relation O on X is called *order* on X if O is reflexive, transitive, and antisymmetric. The order O is called *linear* if for all $x, \tilde{x} \in X$ either $(x, \tilde{x}) \in O$ or $(\tilde{x}, x) \in O$.

All orders discussed in Examples 1.17 are those orders on \mathbb{N} , \mathbb{Z} , and \mathbb{Q} which you are familiar with. In our attempt of presenting a self-contained constructive approach to introduce the real numbers, we include the formal definitions below.

Note that the order on \mathbb{N} which we mention in Examples 1.17.i can be easily defined using elementary set theory.

These definitions are not very enlightening and they will not play a crucial part throughout the remainder of Analysis 1.

³Fields will be defined shortly

EXAMPLES 1.17.

- i. The relation $O_{\mathbb{N}} = \{(n, m) \in \mathbb{N} \times \mathbb{N} : n \leq m\}$ is a linear order on \mathbb{N} .
- ii. The relation $O_{\mathbb{Z}} = \left\{ \left([(n, m)], [(\tilde{n}, \tilde{m})] \right) \in \mathbb{Z} \times \mathbb{Z} : n + \tilde{m} \leq \tilde{n} + m \right\}$ extends the order on \mathbb{N} to the integers \mathbb{Z} .
- iii. The relation $O_{\mathbb{Q}} = \left\{ \left([(z, m)], [(\tilde{z}, \tilde{m})] \right) \in \mathbb{Q} \times \mathbb{Q} : z \cdot \tilde{m} \leq \tilde{z} \cdot m \right\}$ extends the order on \mathbb{Z} to the rational numbers \mathbb{Q} .

In the following we shall simply write $r \leq \tilde{r}$ if $(r, \tilde{r}) \in O_{\mathbb{Q}}$.

DEFINITION 1.18. A field F is called *ordered* if

- (O1) There exists an order ' \leq ' on F .
- (O2) The order is linear, that is, for all $x, y \in F$ either $x < y$ or $x > y$ or $x = y$.
- (O3) $x \leq y$ implies $x + z \leq y + z$ for all $x, y, z \in F$ and if $x, y > 0$ then $x \cdot y > 0$.

DEFINITION 1.19. An ordered field F is called *archimedean* if for all $x, y \in F$, $x, y > 0$ exists $n \in \mathbb{N}$ with

$$nx := \underbrace{x + x + \dots + x}_{n\text{-times}} > y.$$

THEOREM 1.20. The set of rational numbers \mathbb{Q} together with the two binary operations addition and multiplication defined in Examples 1.14.ii and the order given in Examples 1.17.iii is an archimedean ordered field.

1.3. Real numbers

Given a right angled, isosceles triangle with two sides of length 1. What is the length l of the third side?

According to Pythagoras, we have $l^2 = 1^2 + 1^2 = 1 + 1 = 2$. We shall write $l = \sqrt{2}$.

THEOREM 1.21. $\sqrt{2} \notin \mathbb{Q}$, that is, there exists no $m \in \mathbb{Z}$ and $n \in \mathbb{N}$ with $\left(\frac{m}{n}\right)^2 = 2$.

We conclude that there exist line segments with non rational length. Can we define a set $S \supseteq \mathbb{Q}$ containing all “lengths”, and to which we can extend all arithmetic properties of \mathbb{Q} ? Yes, we can!

DEFINITION 1.22. A *Dedekind-cut* $A|B$ in \mathbb{Q} is a pair of subsets A, B of \mathbb{Q} with

- i. $A \cup B = \mathbb{Q}$, $A \neq \emptyset$ and $B \neq \emptyset$, $A \cap B = \emptyset$,
- ii. for all $a \in A$ and $b \in B$ we have $a < b$, that is, $a \leq b$ and $a \neq b$, and
- iii. A contains no largest element.

EXAMPLES 1.23. $\{q \in \mathbb{Q} : q < 2\}|\{q \in \mathbb{Q} : q \geq 2\}$ and $\{q \in \mathbb{Q} : q < 0 \text{ or } q^2 < 2\}|\{q \in \mathbb{Q} : q \geq 0 \text{ and } q^2 \geq 2\}$ are cuts, but $\{q \in \mathbb{Q} : q \leq 2\}|\{q \in \mathbb{Q} : q > 2\}$, $\{q \in \mathbb{Q} : q^2 \leq 2\}|\{q \in \mathbb{Q} : q^2 > 2\}$ and $\{q \in \mathbb{Q} : q < 2\}|\{q \in \mathbb{Q} : q \geq 3\}$ are not.

DEFINITION 1.24. Dedekind-cuts in \mathbb{Q} are called *real numbers*, the set of all real numbers is denoted by \mathbb{R} .

REMARK 1.25. We can embed rational numbers in \mathbb{R} via

$$p \mapsto p^* := \{q \in \mathbb{Q} : q < p\}|\{q \in \mathbb{Q} : q \geq p\}.$$

A cut of the form $p^* := \{q \in \mathbb{Q} : q < p\}|\{q \in \mathbb{Q} : q \geq p\}$, $p \in \mathbb{Q}$ is called rational cut in \mathbb{Q} . The embeddings discussed so far are $\mathbb{N} \hookrightarrow \mathbb{Z} \hookrightarrow \mathbb{Q} \hookrightarrow \mathbb{R}$. Since \hookrightarrow denotes injective maps which respect algebraic properties, we shall omit the $*$ notation and identify elements in the domain with the corresponding elements in the range. That is, we shall write

$$\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R}.$$

At this point of time, we have not defined any algebraic operations on \mathbb{R} (the set of Dedekind cuts in \mathbb{Q}), but we will do this shortly.

DEFINITION 1.26. Let X be a linearly ordered set, $S \subseteq X$. $M \in X$ is an *upper* [resp. *lower*] *bound* of S , if for each $s \in S$ we have $s \leq M$ [resp. $s \geq M$]. If there exists an upper [resp. lower] bound $M \in X$, then we call S *bounded above* [resp. *bounded below*].

$M_0 \in X$ is called the *least upper bound* or *supremum* [resp. *greatest lower bound* or *infimum*] of $S \subseteq X$ if for all upper [lower] bounds $M \in X$ we have $M_0 \leq M$ [resp. $M_0 \geq M$]. The least upper bound [resp. greatest lower bound] of the set S is denoted by $\sup S$ [resp. $\inf S$].

DEFINITION 1.27. (LUP) An ordered set X has the *least upper bound property* if any nonempty subset S of X which is bounded above has a least upper bound (in X).

DEFINITION 1.28. On \mathbb{R} , that is, on the set of Dedekind cuts in \mathbb{Q} , we define:

- i. A linear **order** ' \leq ' on \mathbb{R} via $A|B \leq C|D$ if $A \subseteq C$.
- ii. For $x = A|B$, $y = C|D \in \mathbb{R}$ we set

$$E := \{e \in \mathbb{Q} : \text{there exists } a \in A \text{ and } c \in C \text{ with } e = a + c\}, \quad F := \mathbb{Q} \setminus E$$

and define **addition** on \mathbb{R} via

$$x + y = A|B + C|D := E|F.$$

Further we set $-x = A^-|B^-$, with $A^- = \{-b, b \in B \setminus \{\text{smallest element of } B \text{ (if it exists)}\}\}$ and $B^- = \mathbb{Q} \setminus A^-$.

(Note that $-(-x) = x$, that $x + (-x) = 0^*$ for all $x \in \mathbb{R}$, that $x \geq 0$ if and only if $-x \leq 0$, and that $q^* + \tilde{q}^* = (q + \tilde{q})^*$ and $(-q)^* = -q^*$ for all $q, \tilde{q} \in \mathbb{Q}$.)

- iii. For $x = A|B \geq 0^*$, $y = C|D \geq 0^* \in \mathbb{R}$ we set

$$G := \{e \in \mathbb{Q} : e \leq 0 \text{ or there exists } a > 0 \in A \text{ and } c > 0 \in C \text{ with } e = a \cdot c\}, \quad H := \mathbb{Q} \setminus G$$

and define the **product**

$$x \cdot y = A|B \cdot C|D := G|H.$$

If $x \geq 0$ and $y < 0$ set $x \cdot y = -(x \cdot (-y))$, if $x < 0$ and $y \geq 0$ set $x \cdot y = -((-x) \cdot y)$, and if $x < 0$ and $y < 0$ set $x \cdot y = (-x) \cdot (-y)$. Hence, we have (well) defined **multiplication**

$$\cdot : \mathbb{R} \times \mathbb{R} \longrightarrow \mathbb{R}, \quad (x, y) \mapsto x \cdot y$$

(Note that $q^* \cdot \tilde{q}^* = (q\tilde{q})^*$ for all $q, \tilde{q} \in \mathbb{Q}$.)

THEOREM 1.29. The set of Dedekind cuts in \mathbb{Q} denoted by \mathbb{R} together with the order, the two binary operations addition and multiplication defined above is an archimedean ordered field which satisfies the least upper bound property.

THEOREM 1.30. UNIQUENESS OF THE REAL NUMBER SYSTEM. \mathbb{R} is unique in the following sense: Let F be an archimedean ordered field which has the least upper bound property. Then there exists a bijective mapping $u : F \longrightarrow \mathbb{R}$ which preserves addition, multiplication and order.

Proof. (Sketch) Let F be an archimedean ordered field with the least upper bound property. First note that $1_F >_F 0_F$ since $1_F \neq_F 0_F$ and if $1_F <_F 0_F$ we get $-1_F >_F 0_F$ by (O3) and $1_F = (-1_F)(-1_F) >_F 0_F$ by (O3), a contradiction to (O1). Further, observe that \mathbb{N} can be embedded into F via

$$i : \mathbb{N} \longrightarrow F, \quad n \mapsto n_F = \underbrace{1_F + 1_F + \dots + 1_F}_{n\text{-times}}.$$

By definition we have $n_F + m_F = (n + m)_F$. The injectivity of this mapping follows from an inductive argument using $n_F + 1_F >_F n^* + 0_F$. Let us also note that implies that the order on \mathbb{N} is preserved under i , a very important fact as we shall see later. Further, all $n_F > 0_F$ have an inverse element with respect to addition in F and we may extend i injectively to \mathbb{Z} by setting $n \mapsto -(-n)_F$ for $n < 0$. We can show that $n_F + m_F = (n + m)_F$ still holds, now for all $n, m \in \mathbb{Z}$. Note that (F1) together with (O3) on F implies that $-1_F <_F 0$, since else, we would have $-1_F >_F 0_F$ and $0_F >_F 1_F$.

Further, we can use the same strategy to extend i to cover all rational numbers by setting

$$i : \mathbb{Q} \longrightarrow F, \quad \frac{n}{m} \mapsto \frac{n_F}{m_F} = n_F \cdot m_F^{-1}.$$

(To detail this proof, we would have to show that i is well defined, that is, that the image of q under i does not depend on the particular representation of q as fraction of integer and natural number.)

Note that, again, we have $0 < \frac{n}{m} < \frac{\tilde{n}}{\tilde{m}}$ if and only if $0_F <_F \frac{n_F}{m_F} <_F \frac{\tilde{n}_F}{\tilde{m}_F}$ due to (O3) since else $n_F \cdot_F \tilde{m}_F > \tilde{n}_F \cdot_F m_F$. Further $q_F + r_F = (q + r)_F$ and $q_F \cdot_F r_F = (q \cdot_F r)_F$ holds for all $q, r \in \mathbb{Q}$.

After having observed that any ordered field contains a copy of \mathbb{Q} as an ordered subfield, we can proceed to define the "uniqueness" map u :

$$u : F \longrightarrow \mathbb{R}, \quad x \mapsto A_x | B_x = \{q \in \mathbb{Q} : q_F <_F x\} | \{q \in \mathbb{Q} : q_F \geq_F x\}.$$

It remains to show that u is well defined (are these elements on the right really Dedekind cuts?), it preserves addition, multiplication, and order, and that u is bijective. Note that we still have not used the fact that the order on F is archimedean and that F has the least upper bound property.

So let us first look whether the map is well defined. Clearly $A_x \cap B_x = \emptyset$ and $A_x \cup B_x = \mathbb{Q}$. If $x >_F 0_F$ we have $0 \in A_x$ and $B_x \neq \emptyset$ since the archimedean property implies the existence of $n \in \mathbb{N}$ such that

$$n_F = \underbrace{1_F +_F 1_F +_F \dots +_F 1_F}_{n\text{-times}} > x$$

and therefore $n_F \in B_x$. If $x \leq_F 0_F$ we get $B_x \neq \emptyset$ cheaply and we can use a similar argument as above to show that $A_x \neq \emptyset$.

Transitivity shows that for $a \in A_x$ and $b \in B_x$ we have $a_F < x \leq b_F$ and therefore $a \leq b$.

To show that A_x has no largest element, we need to show the following fact, which we shall repeatedly use not only in this proof.

Claim: Let F be an archimedean ordered field which has the least upper bound property and let $x, y \in F$. If $x < y$, then exists $q \in \mathbb{Q}$ such that $x < q_F < y$.

Proof of the claim: Fix $x, y \in F$ with $x < y$. Then $y - x > 0$ and therefore $(y - x)^{-1} > 0$. Pick $m_F > (y - x)^{-1} > 0$. Set $u = \sup\{n \in \mathbb{Z} : \frac{n_F}{m_F} \leq x\}$. Then $x < \frac{u_F + 1_F}{m_F} < y$, since $\frac{u_F + 1_F}{m_F} > y$ would imply $\frac{u_F + 1_F}{m_F} > y > x \geq \frac{u_F}{m_F}$ and $\frac{1_F}{m_F} = \frac{u_F + 1_F}{m_F} - \frac{u_F}{m_F} > y - x > \frac{1}{m_F}$, a contradiction.

The set A_x has no largest element, since for any q_F , ($q \in \mathbb{Q}$) in A_x we can find \tilde{q}_F , ($\tilde{q} \in \mathbb{Q}$) with $x > \tilde{q}_F > q_F$.

We have shown that $A_x | B_x \in \mathbb{R}$, let us now check surjectivity of u . Let $A | B$ be any cut in \mathbb{Q} . Set $A_F = \{q_F \in F : q \in A\}$ and $x_{A|B} = \sup A_F$ which exists due to the l.u.b. property of F . It is easy to see that $u(x_{A|B}) = A_x | B_x = A | B$.

Injectivity follows from the claim proven above (why?). The mapping u preserves multiplication and addition since it does fulfill these properties on \mathbb{Q} and due to the definition of \mathbb{R} and u . \square

That's it for Dedekind cuts, we are done. From now on, we will think of real numbers as elements on the real line, its elements are denoted with letters such as $x, y, a, b, \alpha, \beta, \dots$

THEOREM 1.31. For every real number $x > 0$ and $n \in \mathbb{N}$ exists exactly one real number $y > 0$ with $y^n = x$. This y is called n -th root of x and is denoted by $x^{\frac{1}{n}}$ or $\sqrt[n]{x}$.

THEOREM 1.32. NESTED INTERVAL PROPERTY.

For $n \in \mathbb{N}$, let $I_n = [a_n, b_n] = \{x \in \mathbb{R} : a_n \leq x \leq b_n\} \subset \mathbb{R}$ be closed intervals with $I_n \supseteq I_{n+1}$ for all $n \in \mathbb{N}$. Then $\bigcap_{n \in \mathbb{N}} I_n \neq \emptyset$.

DEFINITION 1.33. A *sequence* a in a set X is a function $a: \mathbb{N} \longrightarrow X, n \mapsto a(n)$. Note that by convention we shall write a_n instead of $a(n)$, and a is often denoted by $(a_n)_{n \in \mathbb{N}}$ or $\{a_n\}_{n \in \mathbb{N}}$. Do not confuse the sequence $a = (a_n)_{n \in \mathbb{N}} = \{a_n\}_{n \in \mathbb{N}}$ with the set $\{a_n, n \in \mathbb{N}\} = \mathcal{R}_a$.

DEFINITION 1.34. A set X is *countable* if there exists a surjective function (sequence) $a: \mathbb{N} \longrightarrow X, n \mapsto a(n)$.

THEOREM 1.35. If the sets $A_n \subset X, n \in \mathbb{N}$, are countable, then $\bigcup_{n \in \mathbb{N}} A_n$ is countable.

COROLLARY 1.36. \mathbb{Q} is countable.

THEOREM 1.37. The set containing all sequences with values in $\{0, 1, 2, \dots, n\}, n \geq 1$, is not countable.

THEOREM 1.38. \mathbb{R} is not countable.

1.4. Complex numbers

We shall now define the complex number system.

DEFINITION 1.39. The cartesian product $\mathbb{R} \times \mathbb{R}$ together with the binary operations

$$\begin{aligned} + & : (\mathbb{R} \times \mathbb{R}) \times (\mathbb{R} \times \mathbb{R}) \longrightarrow \mathbb{R} \times \mathbb{R}, \quad ((a, b), (c, d)) \mapsto (a + c, b + d) \\ \cdot & : (\mathbb{R} \times \mathbb{R}) \times (\mathbb{R} \times \mathbb{R}) \longrightarrow \mathbb{R} \times \mathbb{R}, \quad ((a, b), (c, d)) \mapsto (ac - bd, ad + bc) \end{aligned}$$

form a field with additive neutral element $(0, 0)$ and multiplicative neutral element $(1, 0)$ which is called the *field of complex numbers*.

THEOREM 1.40. The map $G : \mathbb{R} \longrightarrow \mathbb{C}$, $a \mapsto (a, 0)$ is an embedding of the real numbers into the complex numbers, that is, G is injective and we have for all $a, b \in \mathbb{R}$

$$G(a + b) = G(a) + G(b) \quad \text{and} \quad G(ab) = G(a) \cdot G(b).$$

Hence, we can consider \mathbb{R} as a subfield of \mathbb{C} .

PROPOSITION 1.41. For $i := (0, 1)$, we have $i^2 = (-1, 0)$, and for $a, b \in \mathbb{R}$ we have $G(a) + G(b) \cdot i = (a, b)$. From now on we shall consider \mathbb{R} as a subfield of \mathbb{C} and drop the embedding G in our description of complex numbers. Hence, we shall write $a + bi = (a, b) \in \mathbb{C}$.

DEFINITION 1.42. For $z = a + bi \in \mathbb{C}$ with $a, b \in \mathbb{R}$ we shall call $a = \operatorname{Re}(z) \in \mathbb{R}$ the *real part* of z and $b = \operatorname{Im}(z) \in \mathbb{R}$ the *imaginary part* of z . The *conjugate* of z is $\bar{z} = a - bi$ and the *absolute value* of z is $|z| = \sqrt{a^2 + b^2}$.

PROPOSITION 1.43. For all $z = a + bi$, $w = c + di \in \mathbb{C}$ with $a, b, c, d \in \mathbb{R}$ we have

$$\begin{aligned} \operatorname{Re}(z + w) &= \operatorname{Re}(z) + \operatorname{Re}(w) \\ \operatorname{Im}(z + w) &= \operatorname{Im}(z) + \operatorname{Im}(w) \\ \overline{z + w} &= \bar{z} + \bar{w} \\ \overline{z\bar{w}} &= \bar{z} \bar{\bar{w}} \\ |z| + |w| &\geq |z + w| \\ |z||w| &= |zw| \\ z + \bar{z} &= 2\operatorname{Re}(z) \\ z - \bar{z} &= 2i\operatorname{Im}(z) \\ z\bar{z} &= |z|^2 \\ z^{-1} &= \frac{1}{|z|^2} \bar{z} \\ |\operatorname{Re}(z)| &\leq |z| \\ |\operatorname{Im}(z)| &\leq |z|. \end{aligned}$$

REMARK 1.44. A more geometrical treatise of complex numbers is contained in the homework.

2. CONVERGENCE OF SEQUENCES IN METRIC SPACES AND NUMERIC SERIES

The goal of this section is to discuss real and complex valued sequences and series. Many results concerning real and complex sequences hold in a more general setup, that is, in metric spaces. In order to avoid the repetition of arguments, we shall phrase some results in the metric space setup, nevertheless, at this point of time it might be best to think of only two metric spaces, that is, the spaces of real and complex numbers. In these special cases, the distance between two numbers x and y is $|x - y|$.

2.1. Sequences in metric spaces

DEFINITION 2.1. A set X together with a binary function $d : X \times X \longrightarrow \mathbb{R}$ is a *metric space* with *metric* d if d satisfies

- i. $d(x, \tilde{x}) > 0$ if $x \neq \tilde{x}$ and $d(x, x) = 0$ for all $x \in X$,
- ii. $d(x, \tilde{x}) = d(\tilde{x}, x)$ for all $x, \tilde{x} \in X$,
- iii. $d(x, \tilde{x}) \leq d(x, \tilde{x}) + d(\tilde{x}, \tilde{x})$ for all $x, \tilde{x}, \tilde{x} \in X$.

The function d is called *metric* or *distance function* on the set X and we shall denote a metric space by (X, d) or simply by X if it is well understood which metric d on X is being considered.

EXAMPLES 2.2.

- i. The set of real numbers \mathbb{R} with metric $d_2(x, y) = |x - y|$ is a metric space. If no other metric is explicitly mentioned, we shall always consider \mathbb{R} to be equipped with the *euclidean metric* d_2 .
- ii. The set of complex numbers \mathbb{C} with metric $d_2(x, y) = |x - y| = \sqrt{(\operatorname{Re}(x - y))^2 + (\operatorname{Im}(x - y))^2}$ is a metric space. If no other metric is explicitly mentioned, we shall always consider \mathbb{C} to be equipped with the d_2 metric.
- iii. Given any set X , we can define a metric on X via

$$d_0(x, y) = \begin{cases} 0 & \text{if } x = y; \\ 1 & \text{else} \end{cases} \quad \text{for } x, y \in X.$$

This metric is called *discrete metric* on X .

DEFINITION 2.3. A sequence $(x_n)_{n \in \mathbb{N}}$ in \mathbb{R} is said to *converge* to $x_0 \in \mathbb{R}$ if for all $\varepsilon > 0$ exists $N \in \mathbb{N}$ such that

$$|x_n - x_0| < \varepsilon \quad \text{for all naturals } n \geq N.$$

If $(x_n)_{n \in \mathbb{N}}$ converges to x_0 in \mathbb{R} we write $\lim_{n \rightarrow \infty} x_n = x_0$, or $x_n \xrightarrow{n \rightarrow \infty} x_0$, or simply $x_n \longrightarrow x_0$. The element $x_0 \in \mathbb{R}$ is called *limit* of $(x_n)_{n \in \mathbb{N}}$ in \mathbb{R} .

DEFINITION 2.4. A sequence $(x_n)_{n \in \mathbb{N}}$ in a metric space (X, d) is said to *converge* to $x_0 \in X$ if for all $\varepsilon > 0$ (that is, $\varepsilon \in \mathbb{R}$ with $\varepsilon >_{\mathbb{R}} 0_{\mathbb{R}}$) exists $N \in \mathbb{N}$ such that

$$d(x_n, x_0) < \varepsilon \quad \text{for all naturals } n \geq N.$$

If (x_n) converges to x_0 in (X, d) we write $\lim_{n \rightarrow \infty} x_n = x_0$, or $x_n \xrightarrow{n \rightarrow \infty} x_0$, or simply $x_n \rightarrow x_0$. The element $x_0 \in X$ is called *limit* of (x_n) in (X, d) .

EXAMPLES 2.5.

- i. The sequence $(\frac{1}{n})_{n \in \mathbb{N}}$ in (\mathbb{R}, d_2) converges to $0 \in \mathbb{R}$.
- ii. The sequence $(\frac{1}{n})_{n \in \mathbb{N}}$ in (\mathbb{R}, d_0) does not converge to any $x_0 \in \mathbb{R}$, since for any $x_0 \in \mathbb{R}$ we have $d_0(x_0, x_n) < \frac{1}{2}$ for at most one index $n \in \mathbb{N}$.

PROPOSITION 2.6. A sequence $(z_n)_n$ in \mathbb{C} converges in (\mathbb{C}, d_2) (or simply in \mathbb{C}) if and only if

$$\operatorname{Re}(z_n) \xrightarrow{n \rightarrow \infty} \operatorname{Re}(z_0) \text{ in } \mathbb{R}$$

and

$$\operatorname{Im}(z_n) \xrightarrow{n \rightarrow \infty} \operatorname{Im}(z_0) \text{ in } \mathbb{R}.$$

That is, sequences converge in \mathbb{C} if and only if both, real and imaginary part converge in \mathbb{R} . Therefore, a real valued sequence converges in \mathbb{R} if and only if it converges in \mathbb{C} .

THEOREM 2.7. The limit of a converging sequence in a metric space (X, d) is unique, that is, if $x_n \xrightarrow{n \rightarrow \infty} x_0 \in X$ and $x_n \xrightarrow{n \rightarrow \infty} \tilde{x}_0 \in X$, then $x_0 = \tilde{x}_0$.

DEFINITION 2.8. A subset S in a metric space (X, d) is called *bounded* if there exists $x_0 \in X$ and $M \in \mathbb{R}^+$ such that $d(x_0, x) \leq M$ for all $x \in S$.

A sequence (x_n) is *bounded* in (X, d) if its range $\{x_n : n \in \mathbb{N}\}$ is a bounded set in (X, d) .

THEOREM 2.9. Every converging sequence (x_n) in a metric space (X, d) is bounded.

DEFINITION 2.10. A sequence (x_n) in \mathbb{R} is

- i. *monotonically increasing* if $x_n \leq x_{n+1}$ for all $n \in \mathbb{N}$,
- ii. *strictly monotonically increasing* if $x_n < x_{n+1}$ for all $n \in \mathbb{N}$,
- iii. *monotonically decreasing* if $x_n \geq x_{n+1}$ for all $n \in \mathbb{N}$, and
- iv. *strictly monotonically decreasing* if $x_n > x_{n+1}$ for all $n \in \mathbb{N}$.

A sequence is called *monotone* if it is either monotonically increasing or decreasing.

THEOREM 2.11. Monotonic sequences converge in \mathbb{R} if and only if they are bounded.

THEOREM 2.12. ALGEBRAIC LIMIT THEOREM. If $a_n \xrightarrow{n \rightarrow \infty} a_0$ and $b_n \xrightarrow{n \rightarrow \infty} b_0$ in \mathbb{C} . Then

- i. $(a_n + b_n) \xrightarrow{n \rightarrow \infty} a_0 + b_0$,
- ii. $a_n b_n \xrightarrow{n \rightarrow \infty} a_0 b_0$, and
- iii. $\frac{1}{a_n} \xrightarrow{n \rightarrow \infty} \frac{1}{a_0}$ if $a_0, a_n \neq 0$ for $n \in \mathbb{N}$

THEOREM 2.13. ORDER LIMIT THEOREM. If $a_n \xrightarrow{n \rightarrow \infty} a_0$ and $b_n \xrightarrow{n \rightarrow \infty} b_0$ in \mathbb{C} with $a_n \leq b_n$ for all $n \in \mathbb{N}$, then $a_0 \leq b_0$.

EXAMPLES 2.14.

- i. For $p > 0$ we have $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 0$.
- ii. For $p > 0$ we have $\lim_{n \rightarrow \infty} \sqrt[p]{p} = 1$.
- iii. $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$.
- iv. For $p > 0$ and $\alpha \in \mathbb{R}$ we have $\lim_{n \rightarrow \infty} \frac{n^\alpha}{(1+p)^n} = 0$.
- v. If $x \in \mathbb{C}$ with $|x| < 1$, then $\lim_{n \rightarrow \infty} x^n = 0$.

DEFINITION 2.15. Let (x_n) be a sequence in (X, d) and let $n_1 < n_2 < n_3 < \dots$ be a strictly increasing sequence of natural numbers. Then $(x_{n_k})_{k \in \mathbb{N}}$ is called *subsequence* of (x_n) .

EXAMPLE 2.16. Given the sequence $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$, we have $\frac{1}{2}, \frac{1}{4}, \frac{1}{6}, \dots$ is a subsequence of $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$, but $1, \frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$ and $\frac{1}{2}, 1, \frac{1}{4}, \frac{1}{3}, \frac{1}{6}, \frac{1}{5}, \dots$ are not. In general, $(x_{n_k})_{k \in \mathbb{N}}$ with $x_{n_k} = x_{2k}$ is a subsequence of (x_n) .

THEOREM 2.17. Every subsequence $(s_{n_k})_k$ of a convergent sequence $(s_n)_n$ in (X, d) converges to the same limit as $(s_n)_n$.

EXAMPLE 2.18. The sequence $\frac{1}{2}, \frac{1}{2 + \frac{1}{2}}, \frac{1}{2 + \frac{1}{2 + \frac{1}{2}}}, \frac{1}{2 + \frac{1}{2 + \frac{1}{2 + \frac{1}{2}}}}, \dots$, converges to $\sqrt{2} - 1$ in \mathbb{R} .

THEOREM 2.19. BOLZANO–WEIERSTRASS THEOREM. Every bounded sequence $(s_n)_n$ in \mathbb{R} has a converging subsequence.

2.2. The extended real number system, \limsup and \liminf

DEFINITION 2.20. The *extended real number system* is the linear ordered set $\mathbb{R}^* = \mathbb{R} \cup \{+\infty, -\infty\}$ with $-\infty <_{\mathbb{R}^*} x <_{\mathbb{R}^*} y <_{\mathbb{R}^*} +\infty$ for all $x <_{\mathbb{R}} y$ in \mathbb{R} .

Note that the field structure on \mathbb{R} cannot be extended (in a meaningful way) to \mathbb{R}^* . Nevertheless, it is customary to set

$$\begin{aligned} x + (+\infty) &= +\infty \quad \text{for } x \in \mathbb{R}, \\ x + (-\infty) = x - (+\infty) &= -\infty \quad \text{for } x \in \mathbb{R}, \text{ and} \\ \frac{x}{+\infty} = \frac{x}{-\infty} &= 0 \quad \text{for } x \in \mathbb{R}. \end{aligned}$$

If $x > 0$ we set $x \cdot (+\infty) = +\infty$, $x \cdot (-\infty) = -\infty$, if $x < 0$ then $x \cdot (+\infty) = -\infty$ and $x \cdot (-\infty) = +\infty$.

Further, if for all $M \in \mathbb{R}^+$ there exists $N \in \mathbb{N}$ such that

$$x_n \geq M \quad \text{for all naturals } n \geq N,$$

then we write $\lim_{n \rightarrow \infty} x_n = \infty$, or $x_n \xrightarrow{n \rightarrow \infty} \infty$, or simply $x_n \longrightarrow \infty$. Correspondingly, if for all $M \in \mathbb{R}^+$ there exists $N \in \mathbb{N}$ such that

$$x_n \leq -M \quad \text{for all naturals } n \geq N,$$

then we write $\lim_{n \rightarrow \infty} x_n = -\infty$, or $x_n \xrightarrow{n \rightarrow \infty} -\infty$, or $x_n \longrightarrow -\infty$.

PROPOSITION 2.21. The linearly ordered set \mathbb{R}^* has the least upper bound property. Since in addition every subset of \mathbb{R}^* is bounded above by ∞ , each non-empty subset of \mathbb{R}^* has a least upper bound.

PROPOSITION 2.22. Let (x_n) be a sequence of real numbers. Then

$$E_{(x_n)} = \{x_0 \in \mathbb{R}^* : \text{there exists a subsequence } (x_{n_k}) \text{ of } (x_n) \text{ with } x_{n_k} \xrightarrow{k \rightarrow \infty} x_0\} \subseteq \mathbb{R}^*$$

is not empty.

DEFINITION 2.23. Let (x_n) be a sequence of real numbers. Set

$$E_{(x_n)} = \{x_0 \in \mathbb{R}^* : \text{there exists a subsequence } (x_{n_k}) \text{ of } (x_n) \text{ with } x_{n_k} \xrightarrow{k \rightarrow \infty} x_0\} \subseteq \mathbb{R}^*$$

and define

$$\begin{aligned} \limsup_{n \rightarrow \infty} x_n &= \sup E_{(x_n)} = l.u.b. E_{(x_n)} \in \mathbb{R}^*, \text{ and} \\ \liminf_{n \rightarrow \infty} x_n &= \inf E_{(x_n)} = -l.u.b. (-E_{(x_n)}) \in \mathbb{R}^*. \end{aligned}$$

Any $x_0 \in E_{(x_n)} \cap \mathbb{R}$ is called *limit point* of the real valued sequence (x_n) .

EXAMPLES 2.24.

- i. Choose (x_n) such that $\{x_n, n \in \mathbb{N}\} = \mathbb{Q}$. Then $\limsup_{n \rightarrow \infty} x_n = +\infty$ and $\liminf_{n \rightarrow \infty} x_n = -\infty$.
- ii. Let $x_n = (-1)^n \left(1 + \frac{1}{n}\right)$ for $n \in \mathbb{N}$. Then $\limsup_{n \rightarrow \infty} x_n = +1$ and $\liminf_{n \rightarrow \infty} x_n = -1$.

LEMMA 2.25. Let (x_n) be a sequence in \mathbb{R} and $s \in \mathbb{R}^*$. If $s > \limsup_{n \rightarrow \infty} x_n$, then exists $N \in \mathbb{N}$ such that $x_n \leq s$ for all $n \geq N$. If $s < \liminf_{n \rightarrow \infty} x_n$, then exists $N \in \mathbb{N}$ such that $x_n \geq s$ for all $n \geq N$.

Proof. Fix (x_n) and $s \in \mathbb{R}^*$ with $s > \limsup_{n \rightarrow \infty} x_n$. We shall show that there exists $N \in \mathbb{N}$ such that $x_n \leq s$ for all $n \geq N$. The second assertion follows verbatim.

If $s = \infty$, then $s_n \leq s = \infty$ for all $n \geq 1$.

We have $s > \limsup_{n \rightarrow \infty} x_n \geq -\infty$, and, hence, we can turn our attention to the remaining case $s \in \mathbb{R}$. Suppose that for any $N \in \mathbb{N}$ there exists an index $n_N \in \mathbb{N}$ such that $x_{n_N} > s$. In this case, we can pick n_1 such that $x_{n_1} > s$, then $n_2 > n_1$ with $x_{n_2} > s$, and, inductively $n_{k+1} > n_k$, $k \in \mathbb{N}$.

Since (x_{n_k}) is a subsequence of (x_n) and, therefore, any subsequence of (x_{n_k}) is also a subsequence of (x_n) , we have $E_{(x_{n_k})_k} \subseteq E_{(x_n)_n}$. Pick $y \in E_{(x_{n_k})_k} \neq \emptyset$ and observe that an application of the order limit theorem to subsequences of $(x_{n_k})_k$ implies $y \geq s$ since $x_{n_k} \geq s$ for all $k \in \mathbb{N}$. The fact that $y \in E_{(x_n)_n}$ implies $\limsup_{n \rightarrow \infty} x_n \geq y \geq s > \limsup_{n \rightarrow \infty} x_n$, which is nonsense. Contradiction! \square

THEOREM 2.26. Let (x_n) be a sequence in \mathbb{R} . Then for $x_0 \in \mathbb{R}^*$ we have $\lim_{n \rightarrow \infty} x_n = x_0$ if and only if $\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = x_0$.

Proof. Let us first assume $\lim_{n \rightarrow \infty} x_n = x_0 \in \mathbb{R}^*$. Then $E_{(x_n)_n} = \{x_0\}$ and therefore $\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = x_0$.

Let us now assume $\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = x_0$ with $x_0 \in \mathbb{R}$. Fix $\epsilon > 0$ and use Lemma 2.25 to obtain $N \in \mathbb{N}$ such

$$x_0 - \epsilon < \liminf_{n \rightarrow \infty} x_n - \frac{\epsilon}{2} \leq x_n \leq \limsup_{n \rightarrow \infty} x_n + \frac{\epsilon}{2} < x_0 + \epsilon \quad \text{for all } n \geq N.$$

Since $\epsilon > 0$ was chosen arbitrarily, we have that (x_n) converges and $\lim_{n \rightarrow \infty} x_n = x_0$.

Let us assume $\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = +\infty$. Lemma 2.25 implies that for all $M < \infty$ exists $N \in \mathbb{N}$ with $x_n > M$ for $n \geq N$. This gives $\lim_{n \rightarrow \infty} x_n = \infty$.

The case $\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = -\infty$ can be treated in the same way as the case $\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = +\infty$. \square

2.3. Cauchy sequences and complete metric spaces

DEFINITION 2.27. A sequence (x_n) in a metric space (X, d) is called *Cauchy sequence* if for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $d(x_n, x_m) < \varepsilon$ for all $n, m \geq N$.

PROPOSITION 2.28. Any converging sequence in a metric space is a Cauchy sequence.

PROPOSITION 2.29. Any Cauchy sequence in a metric space is bounded.

DEFINITION 2.30. A metric space (X, d) is called complete if all Cauchy sequences in X converge in X .

REMARK 2.31. Not every metric space is complete. For example, consider the punctured real line $\mathbb{R} \setminus \{0\}$ with $d(x, y) = |x - y|$. The sequence $a_n = \frac{1}{n}$ is Cauchy in $\mathbb{R} \setminus \{0\}$ with $d(x, y) = |x - y|$ since for fixed $\epsilon > 0$ we can pick $N > \frac{1}{\epsilon}$ and get

$$d(x_n, x_m) = |x_n - x_m| = \left| \frac{1}{n} - \frac{1}{m} \right| = \left| \frac{m - n}{mn} \right| < \frac{1}{\max\{n, m\}} \leq \frac{1}{N} < \epsilon$$

for all $n, m \geq N$. Nevertheless, (a_n) does not converge in $\mathbb{R} \setminus \{0\}$, since if it would converge to say $\alpha \in \mathbb{R} \setminus \{0\}$, then it is easy to see that for any $\epsilon > 0$ there would exist some N_ϵ such that

$$|\alpha - 0| \leq |\alpha - x_n| + |0 - x_n| < \epsilon + \epsilon = 2\epsilon.$$

Hence $|\alpha - 0| \leq 2\epsilon$ for all $\epsilon > 0$ and therefore $|\alpha - 0| = 0$ and $\alpha = 0$, a contradiction to $\alpha \in \mathbb{R} \setminus \{0\}$.

PROPOSITION 2.32. Let (X, d) be a metric space and (x_n) be a Cauchy sequence with a converging subsequence, that is there exists (x_{n_k}) with $x_{n_k} \xrightarrow{k \rightarrow \infty} x_0$. Then $x_n \xrightarrow{n \rightarrow \infty} x_0$.

THEOREM 2.33. \mathbb{R} and \mathbb{C} are complete.

2.4. Real and complex series

DEFINITION 2.34. Let (a_n) be a sequence in \mathbb{C} . We call the expression $\sum_{n=1}^{\infty} a_n$ *infinite series* in \mathbb{C} . Further, $S_N = a_1 + a_2 + \dots + a_N = \sum_{n=1}^N a_n$ is called the *N-th partial sum* of $\sum_{n=1}^{\infty} a_n$.

If the sequence $(S_N)_{N \in \mathbb{N}}$ of partial sums converges, we set $\sum_{n=1}^{\infty} a_n = \lim_{N \rightarrow \infty} S_N$. (Be aware of the abuse of notation: $\sum_{n=1}^{\infty} a_n$ denotes a series as well as the limit of its partial sums (in case of convergence)).

EXAMPLE 2.35. Let $a \in \mathbb{C}$ with $|a| < 1$. Then $S_N = \sum_{n=0}^N a^n = \frac{a^{N+1} - 1}{a - 1}$ and $\sum_{n=0}^{\infty} a^n = \frac{1}{1 - a}$.

DEFINITION 2.36. Set $e = \sum_{n=0}^{\infty} \frac{1}{n!} \in \mathbb{R}$.

REMARK 2.37. e is well defined:

$$\begin{aligned} S_N = \sum_{n=0}^N \frac{1}{n!} &= 1 + 1 + \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4} + \dots + \frac{1}{N!} \\ &< 1 + 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^{N-1}} \\ &< 1 + \left(\sum_{n=0}^{\infty} \left(\frac{1}{2} \right)^n \right) = 1 + \frac{1}{1 - \frac{1}{2}} = 3 \end{aligned}$$

Hence (S_n) is bounded. Since (S_N) is also monotone, the sequence of partial sums converges and therefore the series $\sum_{n=0}^{\infty} \frac{1}{n!}$ converges.

THEOREM 2.38. $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^n = e$

THEOREM 2.39. e is irrational.

THEOREM 2.40. CAUCHY CRITERION. The complex series $\sum_{n=1}^{\infty} a_n$ converges in \mathbb{C} if and only if for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that

$$\left| \sum_{n=k}^m a_n \right| < \varepsilon \quad \text{for all } k, m \geq N.$$

PROPOSITION 2.41. If $\sum_{n=1}^{\infty} a_n$ converges in \mathbb{C} then $a_n \xrightarrow{n \rightarrow \infty} 0$.

THEOREM 2.42. DOMINATED CONVERGENCE THEOREM (DCT). Let (a_n) be a sequence in \mathbb{C} .

- i. If there exists a real valued, non-negative sequence (b_n) with $\sum_{n=1}^{\infty} b_n$ converges and $|a_n| \leq b_n$ for all $n \geq N_0, n \in \mathbb{N}$ then $\sum_{n=1}^{\infty} a_n$ converges.
- ii. If $a_n \geq b_n > 0$ for $n \geq N_0, n \in \mathbb{N}$ and if $\sum_{n=1}^{\infty} b_n$ diverges, then $\sum_{n=1}^{\infty} a_n$ diverges.

COROLLARY 2.43. Let (a_n) be a sequence in \mathbb{C} . If $\sum_{n=1}^{\infty} |a_n|$ converges, so does $\sum_{n=1}^{\infty} a_n$.

DEFINITION 2.44. A complex valued series $\sum_{n=1}^{\infty} a_n$ with $\sum_{n=1}^{\infty} |a_n|$ converges, is called *absolutely convergent*.

If $\sum_{n=1}^{\infty} a_n$ converges, but $\sum_{n=1}^{\infty} |a_n|$ does not converge, then we call $\sum_{n=1}^{\infty} a_n$ *conditionally convergent*.

DEFINITION 2.45. Let (c_n) be a sequence of complex numbers and let $\pi : \mathbb{N} \rightarrow \mathbb{N}$ be bijective. Then we call the series $\sum_{n=1}^{\infty} c_{\pi(n)}$ a *rearrangement* of the series $\sum_{n=1}^{\infty} c_n$.

THEOREM 2.46.

- i. If $\sum_{n=1}^{\infty} c_n$ converges absolutely, then any rearrangement $\sum_{n=1}^{\infty} c_{\pi(n)}$ converges absolutely to the same limit, that is $\sum_{n=1}^{\infty} c_{\pi(n)} = \sum_{n=1}^{\infty} c_n$ for any bijective $\pi : \mathbb{N} \rightarrow \mathbb{N}$.
- ii. If $(c_n)_n$ is real and if $\sum_{n=1}^{\infty} c_n$ converges conditionally, then for any $x \in \mathbb{R}$ exists bijective $\pi_x : \mathbb{N} \rightarrow \mathbb{N}$ such that $\sum_{n=1}^{\infty} c_{\pi_x(n)} = x$.

EXAMPLE 2.47. Take $S = \sum_{n=1}^{\infty} (-1)^n \frac{1}{n} \neq 0$. Consider:

$$\begin{array}{rcl}
 S & = & -1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \frac{1}{5} + \frac{1}{6} - \frac{1}{7} + \frac{1}{8} - \dots < \frac{1}{2} \\
 + \frac{1}{2}S & = & \quad -\frac{1}{2} \quad \quad + \frac{1}{4} \quad \quad - \frac{1}{6} + \dots + \frac{1}{8} \\
 \hline
 = \frac{3}{2}S & = & -1 + 0 - \frac{1}{3} + \frac{1}{2} - \frac{1}{5} + 0 - \frac{1}{7} + \frac{1}{4} + \dots \\
 \text{but } \frac{3}{2}S & \neq & -1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \frac{1}{5} + \frac{1}{6} - \frac{1}{7} + \frac{1}{8} - \dots = S
 \end{array}$$

since $S \neq 0$. Hence, $\sum_{n=1}^{\infty} (-1)^n \frac{1}{n}$ converges conditionally.

The following criterion is helpful to prove convergence of series which do not converge absolutely.

THEOREM 2.48. LEIBNIZ CRITERION FOR ALTERNATING SERIES. Let (a_n) be a decreasing sequence of positive real numbers with $a_n \rightarrow 0$. Then $\sum_{n=1}^{\infty} (-1)^n a_n$ converges.

THEOREM 2.49. CAUCHY CONDENSATION THEOREM. Suppose $a_1 \geq a_2 \geq \dots \geq 0$. Then $\sum_{n=1}^{\infty} a_n$ converges if and only if $\sum_{k=1}^{\infty} 2^k a_{2^k}$ converges.

PROPOSITION 2.50. For $p \in \mathbb{R}$ we have $\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges if and only if $p > 1$.

THEOREM 2.51. ROOT TEST. Given a complex series $\sum a_n$, set $\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$.

- i. If $\alpha < 1$, then $\sum a_n$ converges absolutely.
- ii. If $\alpha > 1$, then $\sum a_n$ diverges.
- iii. If $\alpha = 1$, then $\sum a_n$ might converge or diverge.

Proof. Here, we shall only show iii.

We have $\limsup_{n \rightarrow \infty} \sqrt[n]{\frac{1}{n}} = \frac{1}{\lim_{n \rightarrow \infty} \sqrt[n]{n}} = 1$ but $\sum \frac{1}{n}$ does not converge.

On the other hand, $\limsup_{n \rightarrow \infty} \sqrt[n]{\frac{1}{n^2}} = 1$ and $\sum \frac{1}{n^2}$ does converge. □

THEOREM 2.52. RATIO TEST. Let $\sum_{n=1}^{\infty} a_n$ be a series of complex numbers.

- i. If $\limsup_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$, then $\sum_{n=1}^{\infty} a_n$ converges absolutely.
- ii. If there exists $N \in \mathbb{N}$ with $\left| \frac{a_{n+1}}{a_n} \right| \geq 1$ for all $n > N$, then $\sum_{n=1}^{\infty} a_n$ diverges.

EXAMPLES 2.53.

- i. Let $a_n = \frac{1}{n}$. Then $\limsup_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \limsup_{n \rightarrow \infty} \frac{n}{n+1} = 1$, but the series $\sum_{n=1}^{\infty} \frac{1}{n}$ does not converge.
- ii. Let $b_n = \frac{1}{n^2}$. Then $\limsup_{n \rightarrow \infty} \frac{b_{n+1}}{b_n} = \limsup_{n \rightarrow \infty} \frac{n^2}{(n+1)^2} = 1$ and $\sum_{n=1}^{\infty} \frac{1}{n^2}$ does converge.

DEFINITION 2.54. The series $\sum_{n=0}^{\infty} c_n z^n$ is called a *power series* with coefficients $c_n \in \mathbb{C}$, $n \in \mathbb{N}$. For $\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{|c_n|} \in [0, \infty] \subset \mathbb{R}^*$ we call

$$R_{(c_n)} = \begin{cases} \frac{1}{\alpha} & \text{if } \alpha \in (0, \infty); \\ \infty & \text{if } \alpha = 0; \\ 0 & \text{if } \alpha = \infty \end{cases}$$

the *radius of convergence* of the power series $\sum_{n=0}^{\infty} c_n z^n$.

THEOREM 2.55. The series $\sum_{n=0}^{\infty} c_n z^n$ converges if $|z| < R_{(c_n)}$ and diverges if $|z| > R_{(c_n)}$, and $\sum_{n=0}^{\infty} c_n z^n$ may or may not converge for $z \in \mathbb{C}$ with $|z| = R_{(c_n)}$.

REMARK 2.56. It is easy to see that a series of the form $\sum_{n=0}^{\infty} c_n (z - z_0)^n$ converges if $|z - z_0| < R_{(c_n)}$ and diverges if $|z - z_0| > R_{(c_n)}$, a fact which is relevant when discussing Taylor series of a function f at a point $z_0 \in \mathbb{R}$. (See Section 4.)

We conclude this section with a brief discussion of the exponential function $\exp(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$, $z \in \mathbb{C}$. To derive the functional equation $\exp(z+w) = \exp(z)\exp(w)$ we use theorem discussing the product of two series. This theorem is based on a diagonal summation of the product:

$$\begin{aligned} (a_0 + a_1 + a_2 + \dots) \cdot (b_0 + b_1 + b_2 + \dots) = & \begin{array}{cccccc} a_0 b_0 & + & a_0 b_1 & + & a_0 b_2 & + & a_0 b_3 & + & \dots \\ + & a_1 b_0 & + & a_1 b_1 & + & a_1 b_2 & + & a_1 b_3 & + & \dots \\ + & a_2 b_0 & + & a_2 b_1 & + & a_2 b_2 & + & a_2 b_3 & + & \dots \\ + & a_3 b_0 & + & a_3 b_1 & + & a_3 b_2 & + & a_3 b_3 & + & \dots \\ & \vdots & & \vdots & & \vdots & & \vdots & & \end{array} \end{aligned}$$

THEOREM 2.57. PRODUCT OF SERIES. Let (a_n) and (b_n) be complex sequences with $\sum_{n=0}^{\infty} a_n = A$ converges absolutely, and $\sum_{n=0}^{\infty} b_n = B$. For $c_n = \sum_{k=0}^n a_k b_{n-k}$, $n \in \mathbb{N}_0$ we have $\sum_{n=0}^{\infty} c_n = A \cdot B$.

COROLLARY 2.58. For $z, w \in \mathbb{C}$ we have $\exp(z + w) = \exp(z) \exp(w)$.

COROLLARY 2.59. For $x \in \mathbb{R}$ we have $\exp(x) = e^x$.

Motivated by this corollary, we shall write e^z for $\exp(z)$ for any $z \in \mathbb{C}$.

3. TOPOLOGY AND CONTINUITY

3.1. Continuous functions

DEFINITION 3.1. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is *continuous* at $x_0 \in \mathbb{R}$ if for all $\varepsilon > 0$ exists $\delta > 0$ s.t. $|f(x) - f(x_0)| < \varepsilon$ if $|x - x_0| < \delta$.

EXAMPLE 3.2. The function

$$f : \mathbb{R} \longrightarrow \mathbb{R}, x \mapsto \begin{cases} x + 2, & \text{if } x \leq -1; \\ x^2, & \text{if } -1 < x < 2; \\ -x + 7, & \text{if } 2 \leq x. \end{cases}$$

is continuous at any point x_0 in $\mathbb{R} \setminus \{2\}$ and discontinuous at $x_0 = 2$.

REMARK 3.3. Continuous functions have some remarkable properties. Most prominently, the intermediate value theorem and the maximum value theorem for real valued functions defined on \mathbb{R} state that given a continuous function $f : [a, b] \rightarrow \mathbb{R}$ then exists $c, d \in \mathbb{R}$, such that $f([a, b]) = [c, d]$. (See Corollary 3.61.)

This theorem can be generalized to metric spaces: If X is a *compact* and *connected* metric space, and $f : X \rightarrow Y$ is *continuous*, then $f(X)$ is *compact* and *connected*. In case of $Y = \mathbb{R}$ we get immediately $f(X) = [c, d]$ for some $c, d \in \mathbb{R}$ since closed intervals are the only subsets of \mathbb{R} which are both, *compact* and *connected*. Well, we need some new vocabulary.

DEFINITION 3.4. Let (X, d_X) and (Y, d_Y) be metric spaces. A function $f : X \rightarrow Y$ is *continuous* at $x_0 \in X$, if for all $\varepsilon \in \mathbb{R} > 0$ exists $\delta > 0$ s.t. $d_Y(f(x), f(x_0)) < \varepsilon$ if $d_X(x_0, x) < \delta$.

DEFINITION 3.5. Let (X, d_X) be a metric space, $x_0 \in X$, and $r \in \mathbb{R}^+$. The *open* [respectively *closed*] *ball* in X of center x_0 and radius r is the set

$$\begin{aligned} B_r(x_0) &= \{x \in X : d_X(x_0, x) < r\} \subseteq X \\ [\text{resp. } B_r^{\text{closed}} &= \{x \in X : d_X(x_0, x) \leq r\}] \end{aligned}$$

We shall also refer to the open ball $B_r(x_0)$ as *r-neighborhood* of x_0 .

THEOREM 3.6. Let (X, d_X) and (Y, d_Y) be metric spaces, and let $f : X \rightarrow Y$, and $x_0 \in X$. f is continuous at x_0 if and only if for all $\varepsilon > 0$ exists $\delta > 0$ s.t. $f(B_\delta(x_0)) \subseteq B_\varepsilon(f(x_0))$.

THEOREM 3.7. CAUCHY–SCHWARZ INEQUALITY.

Let $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{C}$. Then

$$\left| \sum_{i=1}^n a_i \overline{b_i} \right|^2 \leq \sum_{i=1}^n |a_i|^2 \sum_{i=1}^n |b_i|^2$$

EXAMPLES 3.8. Examples of metrics d_0, d_1, d_2, d_∞ on \mathbb{R}^n . Describe respective balls.

THEOREM 3.9. If $f : (\mathbb{R}^n, d_{i_0}) \rightarrow (\mathbb{R}^m, d_{j_0})$ is continuous at $x_0 \in \mathbb{R}^n$ for some $i_0, j_0 \in \{1, 2, \infty\}$, then $f : (\mathbb{R}^n, d_i) \rightarrow (\mathbb{R}^m, d_j)$ for any $i, j \in \{1, 2, \infty\}$.

REMARK 3.10. Obviously, continuity does depend on the metric of choice. Nevertheless, different metrics (not all) lead to the same concept of continuity. We shall now extract the essence of continuous functions between metric spaces which will lead to a whole new class of spaces, namely topological spaces.

DEFINITION 3.11. Let (X, d) be a metric space. $U \subseteq X$ is called (*metric-*) *open* if for each $x_0 \in U$ exists $\varepsilon > 0$ s.t. $B_\varepsilon(x_0) \subseteq U$. A set $A \subseteq X$ is called (*metric-*) *closed* if its complement A^c is (*metric-*) open.

We should check consistency of our vocabulary. We did define *open balls* before defining *open sets*.

THEOREM 3.12. Let (X, d) be a metric space, then open balls are (*metric-*) open.

PROPOSITION 3.13. U is open in (\mathbb{R}^n, d_∞) if and only if U is open in (\mathbb{R}^n, d_1) if and only if U is open in (\mathbb{R}^n, d_2) .

THEOREM 3.14. $f : (X, d_X) \rightarrow (Y, d_Y)$ is continuous on X if and only if $f^{-1}(U)$ is open in (X, d_X) for all U open in (Y, d_Y) .

THEOREM 3.15. Let $\{U_i, i \in I\}$ be a family of (*metric-*) open sets in (X, d) . Then

- i. $U_i \cap U_j$ is open in (X, d) for any $i, j \in I$,
- ii. $\bigcup_{i \in I} U_i$ is open in (X, d) , and
- iii. \emptyset, X are open.

Let us now provide a very important and useful result for the understanding of open sets in subspaces of metric spaces. This result will be used extensively when discussing connected subsets of metric spaces.

THEOREM 3.16. INHERITANCE PRINCIPLE. Let (X, d_X) be a metric space and $A \subseteq X$. Then (A, d_A) becomes a metric space when setting $d_A = d_X|_{A \times A}$, that is, $d_A(a, b) = d_X(a, b)$ for $a, b \in A$. Further, the following hold:

- i. $B \subset A$ is open in (A, d_A) if and only there exists \tilde{B} open in (X, d_X) such that $B = A \cap \tilde{B}$.
- ii. $B \subset A$ is closed in (A, d_A) if and only there exists \tilde{B} closed in (X, d_X) such that $B = A \cap \tilde{B}$.
- iii. $B \subset A$ is clopen (closed and open) in (A, d_A) **if** there exists \tilde{B} clopen in (X, d_X) such that $B = A \cap \tilde{B}$.

3.2. Topological spaces

Theorem 3.15 provides all properties of metric spaces needed to extend the concept of continuous maps on metric spaces to maps between more general spaces. We shall use these properties to define topological spaces.

DEFINITION 3.17. Let X be any set and let \mathcal{T} be a collection of subsets of X with

- i. $\bigcup_{U \in \mathcal{S}} U \in \mathcal{T}$ if $\mathcal{S} \subseteq \mathcal{T}$
- ii. $\bigcap_{U \in \mathcal{S}} U \in \mathcal{T}$ if $\mathcal{S} \subseteq \mathcal{T}$ with \mathcal{S} is a finite set
- iii. $X, \emptyset \in \mathcal{T}$

Then we call \mathcal{T} a *topology* on the *topological space* X , the members U of \mathcal{T} are called (*topology-*) *open*.

EXAMPLE 3.18.

- i. Any set X becomes a topological space when choosing the trivial topology $\mathcal{T} = \{\emptyset, X\}$. This topology is also called *indiscrete* topology.
- ii. Any set X becomes a topological space when choosing as topology the powerset of X , that is, $\mathcal{T} = \mathcal{P}(X)$. This topology is also called *discrete* topology.
- iii. The metric open sets in a metric space (X, d) form a topology on X (see Theorem 3.15). This topology is *induced* by the metric d and we denote it by \mathcal{T}_d .
- iv. Note that for any set X and discrete metric d_0 on X , (ii) and (iii) lead to the same topology, that is, $\mathcal{T}_{d_0} = \mathcal{P}(X)$. This is easy to see since in (X, d_0) (d_0 denotes the discrete metric) we have that $B_1(x) = \{x\}$ for any $x \in X$. Hence, all singletons (sets with only one element) are open and any $S \in \mathcal{P}(X)$ is open since it can be written as union of open sets, for example, $S = \bigcup_{x \in S} \{x\}$.

REMARK 3.19. Recall that, using those properties of (metric-) open sets in a metric space (X, d) that the concept of continuity is based on, we introduced a new family of spaces which is custom made to study continuous maps.

Many properties of metric induced topologies now serve as defining properties when dealing with general topological spaces. E.g., given a topological space (X, \mathcal{T}) and a subset A in X , we can equip A with the so called relative topology $\mathcal{T}_A = \{A \cap U : U \in \mathcal{T}\}$ to obtain a topological space (A, \mathcal{T}_A) . (Compare to the inheritance principle, Theorem 3.16.)

By virtue of Theorem 3.14 we can extend the concept of continuous maps to general topological spaces:

DEFINITION 3.20. Let (X, \mathcal{T}) , (Y, \mathcal{F}) be topological spaces. A function $f : X \rightarrow Y$ is called *continuous* if $f^{-1}(V) \in \mathcal{T}$ for all $V \in \mathcal{F}$.

THEOREM 3.21. Let (X, \mathcal{T}) , (Y, \mathcal{F}) , and (Z, \mathcal{S}) be topological spaces and $f : X \longrightarrow Y$ and $g : Y \longrightarrow Z$ be continuous. Then $g \circ f : X \longrightarrow Z$, $x \mapsto g \circ f(x) = g(f(x))$ is continuous.

Proof. For $U \in \mathcal{S}$ we have $g^{-1}(U) \in \mathcal{F}$ since g is continuous and $(g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U)) \in \mathcal{T}$ since f is continuous. Hence $g \circ f$ continuous \square

In the mathematical discipline topology, one studies whether two topological spaces X and Y have “identical topologies”, that is, whether there exists a continuous, bijective map which maps open sets to open sets (that is, f^{-1} (which exists and is defined on all of Y since f is bijective) is continuous as well).

DEFINITION 3.22. If $f : X \longrightarrow Y$ is bijective and continuous, and if the function $f^{-1} : Y \longrightarrow X$ is continuous as well then we call f a *homeomorphism*.

DEFINITION 3.23. The topological spaces (X, \mathcal{T}) and (Y, \mathcal{F}) are called *homeomorph* if there exists a homeomorphism $f : X \longrightarrow Y$.

DEFINITION 3.24. A sequence (x_n) in the topological space (X, \mathcal{T}) *converges* to x_0 in (X, \mathcal{T}) , if for all $U \in \mathcal{T}$ with $x_0 \in U$ there exists $N_0 \in \mathbb{N}$ s.t. $x_n \in U$ if $n \geq N_0$.

Our back is covered:

THEOREM 3.25. A sequence (x_n) converges to x_0 in the metric space (X, d) if and only if x_n converges to x_0 in the topological space (X, \mathcal{T}_d) .

EXAMPLE 3.26. The function

$$f : [0, 2\pi) \longrightarrow \mathcal{R}_f = \{z \in \mathbb{C} : |z| = 1\} \subset \mathbb{C}, \quad x \mapsto \cos(x) + i \sin(x)$$

is continuous, 1-1, surjective, and continuous, but f^{-1} is not continuous at $1 = \cos(0) + i \sin(0)$. Hence, f is not a homeomorphism. (We shall define \cos and \sin in Section 4.3. At this point of time, we only assume High-School knowledge of trigonometric functions.)

To see this, observe that $\lim_{n \rightarrow \infty} \cos(2\pi - \frac{1}{n}) + i \sin(2\pi - \frac{1}{n}) = 1$, but its image under f^{-1} is the sequence $(f^{-1}(\cos(2\pi - \frac{1}{n}) + i \sin(2\pi - \frac{1}{n})))_n = (2\pi - \frac{1}{n})_n$ which does not converge in $[0, 2\pi)$

In fact, we shall see later that $[0, 2\pi)$ and $\mathcal{R}_f = \{z \in \mathbb{C} : |z| = 1\}$ are not homeomorphic, that is, there exist no homeomorphism $f : [0, 2\pi) \longrightarrow \{z \in \mathbb{C} : |z| = 1\}$.

EXAMPLE 3.27. In the following table we shall consider sequences in \mathbb{R} where \mathbb{R} is equipped with different topologies.

	$\mathcal{T}_{d_0} = \mathcal{P}(\mathbb{R})$	$\mathcal{T} = \{\emptyset, \mathbb{R}\}$	\mathcal{T}_{d_2}
$x_n = 1, \forall n \in \mathbb{N}$	$\lim_{n \rightarrow \infty} x_n = 1$	$\lim_{n \rightarrow \infty} x_n = x$ for any $x \in \mathbb{R}$	$\lim_{n \rightarrow \infty} x_n = 1$
$y_n = \frac{1}{n}, \forall n \in \mathbb{N}$	(y_n) does not converge	$\lim_{n \rightarrow \infty} y_n = y$ for any $y \in \mathbb{R}$	$\lim_{n \rightarrow \infty} y_n = 0$
$z_n = n, \forall n \in \mathbb{N}$	(z_n) does not converge	$\lim_{n \rightarrow \infty} z_n = z$ for any $z \in \mathbb{R}$	(z_n) does not converge
$u_n = (1 + \frac{1}{n})^n$	(u_n) does not converge	$\lim_{n \rightarrow \infty} u_n = u$ for any $u \in \mathbb{R}$	$\lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n = e$

The ambivalence in column $\mathcal{T} = \{\emptyset, \mathbb{R}\}$ is only possible since the topology is not induced by a metric on \mathbb{R} . (We have shown earlier that a sequence in a metric space can only converge to one point.)

DEFINITION 3.28. A subset A of a topological space (X, \mathcal{T}) is called *closed* if $A^C = X \setminus A \in \mathcal{T}$, that is if A^C , the complement of A , is open.

THEOREM 3.29. Let (X, d) be a metric space, then A is closed in (X, \mathcal{T}_d) if and only if given any sequence (x_n) in A with $x_n \rightarrow x_0 \in X$ then automatically $x_0 \in A$.

REMARK 3.30. The characterization of closed sets in metric spaces in Theorem 3.29 does not hold in general topological space.

Continuity at a point $x_0 \in X$ can be described in numerous ways.

THEOREM 3.31. Let $(X, d_X), (Y, d_Y)$ be metric spaces, $x_0 \in X$, and $f : X \rightarrow Y$. The following are equivalent:

- i. The function f is continuous at x_0 , that is, for all $\varepsilon > 0$ exists some $\delta > 0$ such that $d(x_0, x) < \delta$ implies $d(f(x_0), f(x)) < \varepsilon$.
- ii. For all $\varepsilon > 0$ exists some $\delta > 0$ such that $f(B_\delta(x_0)) \subseteq B_\varepsilon(f(x_0))$.
- iii. For all sequences (x_n) in X with $\lim_{n \rightarrow \infty} x_n = x_0$ we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.
- iv. For all open sets U in Y with $x_0 \in U$ exists V open in X with $f(V) \subseteq U$.

THEOREM 3.32. Let $(X, d_X), (Y, d_Y)$ be metric spaces and $f : X \rightarrow Y$. The following are equivalent:

- i. The function f is continuous on X , that is for all $x_0 \in X$ and for all $\varepsilon > 0$ exists some $\delta > 0$ such that $d(x_0, x) < \delta$ implies $d(f(x_0), f(x)) < \varepsilon$.
- ii. For all $x_0 \in X$ and for all $\varepsilon > 0$ exists some $\delta > 0$ such that $f(B_\delta(x_0)) \subseteq B_\varepsilon(f(x_0))$.
- iii. For all $x_0 \in X$ and for all sequences (x_n) in X with $\lim_{n \rightarrow \infty} x_n = x_0$ we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.
- iv. For all open sets U in Y we have $f^{-1}(U)$ is open in X .
- v. For all closed sets A in Y we have $f^{-1}(A)$ is closed in X .

DEFINITION 3.33. Let (X, \mathcal{T}) be a topological space and let $A \subseteq X$.

- i. The *interior* A° of A is given by $A^\circ = \bigcup_{\substack{U \subseteq A \\ U \in \mathcal{T}}} U$.
- ii. The *closure* \bar{A} of A is given by $\bar{A} = \bigcap_{\substack{C \supseteq A \\ C \text{ closed}}} C$.
- iii. The *boundary* ∂A of A is given by $\partial A = \bar{A} \cap \overline{A^C}$.
- iv. A' denotes the *set of all cluster points*, that is $A' = \{x_0 \in X \text{ s.t. there exists a sequence } (x_n) \text{ in } A \text{ with } \lim_{n \rightarrow \infty} x_n = x_0 \text{ and } x_n \neq x_0\}$.

3.3. Compactness

Even though the concept of compact and connected sets and spaces are of topological nature, we shall restrict our treatise to metric spaces (which certainly are just a special breed of topological spaces.)

DEFINITION 3.34. Let A be a subset of a metric space (X, d) and let \mathcal{U} and \mathcal{V} be collections of subsets of X .

- i. The family \mathcal{U} is a *covering of A* if $A \subseteq \bigcup_{U \in \mathcal{U}} U$.
- ii. The family \mathcal{V} is a \mathcal{U} -*subcovering of A* if $\mathcal{V} \subseteq \mathcal{U}$ and $A \subseteq \bigcup_{U \in \mathcal{V}} U$.
- iii. A family of sets \mathcal{U} is called *open* if all $U \in \mathcal{U}$ are open
- iv. The family \mathcal{U} is *finite* if \mathcal{U} consists of finitely many sets (which in turn might contain infinitely many elements of X .)

DEFINITION 3.35. A subset A of a metric space (X, d) is called (*covering-*) *compact* if **every** open cover \mathcal{U} of A contains a finite \mathcal{U} -subcover \mathcal{V} .

EXAMPLES 3.36.

- i. Any finite set is compact.
- ii. The set $\{\frac{1}{n} : n \in \mathbb{N}\}$ is not compact.
- iii. The set $\{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$ is compact.
- iv. In general, let (x_n) be a converging sequence in the metric space (X, d) . Then $\{x_n : n \in \mathbb{N}\} \cup \{\lim_{n \rightarrow \infty} x_n\}$ is compact.
- v. The open interval $(0, 1) \subset \mathbb{R}$ is not compact in (\mathbb{R}, d_2) , since $\mathcal{U} = \{(\frac{1}{n}, 1)\}$ is an open cover of $(0, 1)$ which contains no finite \mathcal{U} -subcover.

DEFINITION 3.37. A subset A in the metric space (X, d) is *sequentially compact* if any sequence (a_n) in A has a subsequence (a_{n_k}) with $\lim_{k \rightarrow \infty} a_{n_k} = a_0$ and $a_0 \in A$.

One of the main goals of this section is to prove that in metric spaces sequentially compactness and covering compactness are the same, that is, a set A is sequentially compact if and only if A is covering compact. Be aware that this theorem does not hold in general topological spaces.

Before proving this theorem, we shall discuss some consequences of compactness.

THEOREM 3.38. Let (X, d) be a metric space and $A \subseteq X$ be compact. If $B \subset A$ is closed in X , then B is compact. Shortly: closed subsets of compact sets are compact.

THEOREM 3.39. Any compact set A in (X, d) is bounded, that is, compact sets are bounded.

THEOREM 3.40. Any infinite subset B of a compact set A in (X, d) has at least one cluster point in A .

THEOREM 3.41. Any compact set is closed.

Theorem 3.41 combines with Theorem 3.39 to the statement that compact sets are closed and bounded. Does the converse hold? It would be nice, we would get a criterium for compactness which is easy to check. Sadly, the converse does not hold in general (see Remark 3.49, but it does hold in euclidean space, that is, \mathbb{R}^n).

To prove the main result of this chapter, we need to introduce the concept of a Lebesgue number.

DEFINITION 3.42. Let \mathcal{U} be a covering of a set A in the metric space (X, d) . Any number $\lambda > 0$ with the property that for all $a \in A$ exists $U \in \mathcal{U}$ such that $B_\lambda(a) \subseteq U$ is called *Lebesgue number* for the covering \mathcal{U} of A .

LEMMA 3.43. Let \mathcal{U} be an open covering of a sequentially compact set A in the metric space (X, d) . Then exists a Lebesgue number $\lambda > 0$ for the covering \mathcal{U} of A .

Proof. Assume there is an open cover $\mathcal{U} = (U_i)_{i \in I}$ of X without a Lebesgue-number, that is for all $n \in \mathbb{N}$ we can choose some $x_n \in X$ such that for all $B_{\frac{1}{n}}(x_n) \not\subseteq U_i$ for all $i \in I$.

Since A is sequential compact, we can extract a convergent subsequence $(x_{n_k})_k$ of (x_n) and set $x_0 := \lim_k x_{n_k} \in X$. Since \mathcal{U} is a covering, we have $x_0 \in U_{i_0}$ for some $i_0 \in I$. Since U_{i_0} is open, there is an $n \in \mathbb{N}$ such that $B_{\frac{1}{n}}(x_0) \subseteq U_{i_0}$.

Pick $K \in \mathbb{N}$ such that $K \geq 2n$ and $d(x_{n_K}, x_0) < \frac{1}{2n}$. We have $B_{\frac{1}{n_K}}(x_{n_K}) \subseteq B_{\frac{1}{n}}(x_0)$ since $d(x, x_{n_K}) < \frac{1}{2n}$ implies $d(x_0, x) < d(x_0, x_{n_K}) + d(x_{n_K}, x) < \frac{1}{2n} + \frac{1}{2n} = \frac{1}{n}$.

We conclude that $B_{\frac{1}{n_K}}(x_{n_K}) \subseteq B_{1/n}(x_0) \subseteq U_{i_0}$, a contradiction. \square

Now we shall provide the main result of this chapter.

THEOREM 3.44. Let (X, d) be a metric space. A set $A \subseteq X$ is sequentially compact if and only if it is covering compact.

Proof. Suppose A is covering compact. Let (x_n) be an arbitrary sequence in A . We have to find a convergent subsequence.

Cover A with balls of radius 1. Since (by covering-compactness) finitely many of them suffice, we throw away all but finitely many of them. Now among the remaining finitely many balls there has to be at least one ball containing x_n for infinitely many values of n . Let us call this ball B_1 . Let n_1 be an index such that x_{n_1} is contained in B_1 .

Now we do the same thing again: cover the set $\overline{B_1} \cap A$, which is a covering-compact set, with (finitely many!) balls of radius $\frac{1}{2}$; one of them, which we call B_2 , must have the property that $B_2 \cap B_1$ is visited infinitely often by the sequence. Choose $n_2 > n_1$ such that $x_{n_2} \in B_2 \cap B_1$. Now continue with $\overline{B_2}$ and radius $\frac{1}{4}$ to construct B_3 and n_3 and continue the process.

Set $C_n = \overline{\bigcap_{k=1}^n B_k} \cap A$ and observe that sequence $X \supseteq C_1 \supseteq C_2 \supseteq \dots$. Since the nested intersection of compact sets whose diameter tends to zero is a single point $x_0 \in A$ (check!), we get by construction, $x_{n_k} \rightarrow x_0$. Since A is closed, we have $x \in A$.

Let us now suppose that A is sequentially compact. Let $\mathcal{U} = (U_i)_{i \in I}$ be an arbitrary open cover. We want to show that \mathcal{U} admits a finite subcover. By Lemma 3.43, this cover has a Lebesgue-number $\lambda > 0$: every $x \in X$ has an $i = i(x)$ such that $B_\lambda(x) \subseteq U_{i(x)}$.

Choose any $x_1 \in X$. Then either $U_1 := U_{i(x_1)}$ covers X and we are done. Otherwise choose any $x_2 \in X \setminus U_1$ and set $U_2 := U_{i(x_2)}$. Again, either $U_1 \cup U_2$ already covers X and we are done, or we can choose $x_3 \in X \setminus (U_1 \cup U_2)$ and so on. Either X is covered after a finite number of steps, or this construction produces an infinite sequence (x_n) in X . However, this sequence has no convergent subsequence, because for all $m \neq n$, $d(x_m, x_n) \geq \lambda$. Hence this case is impossible. \square

LEMMA 3.45. For $a \leq b$ we have $[a, b]$ is compact in \mathbb{R} . (Recall, if not specified we let $d = d_2$ in \mathbb{R}^n .)

LEMMA 3.46. Let A be compact in (\mathbb{R}^n, d_i) and B be compact in (\mathbb{R}^m, d_j) , $i, j \in \{1, 2, \infty\}$. Then $A \times B$ is compact in (\mathbb{R}^{n+m}, d_k) , $k = 1, 2, \infty$.

Proof. Since the topology on (\mathbb{R}^n, d_i) , (\mathbb{R}^m, d_j) and (\mathbb{R}^{n+m}, d_k) does not depend on $i, j, k \in \{1, 2, \infty\}$, we may assume that $i = j = k = 1$.

For $((x_n, y_n))_{n \in \mathbb{N}}$ we have $\lim_{n \rightarrow \infty} (x_n, y_n) = (x_0, y_0)$ in (\mathbb{R}^{n+m}, d_1) if and only if $\lim_{n \rightarrow \infty} x_n = x_0$ in (\mathbb{R}^n, d_1) and $\lim_{n \rightarrow \infty} y_n = y_0$ in (\mathbb{R}^m, d_1) , since $d_1((x_n, y_n), (x_0, y_0)) = d_1(x_n, x_0) + d_1(y_n, y_0)$.

Let $((a_n, b_n))_{n \in \mathbb{N}}$ be a sequence in $A \times B$. We shall construct a subsequence of $((a_n, b_n))_{n \in \mathbb{N}}$ which converges in $A \times B$.

Using sequential compactness of A , we choose a subsequence $(a_{n_k})_{k \in \mathbb{N}}$ of $(a_n)_{n \in \mathbb{N}}$ which converges to $a_0 \in A$. Similarly, we pick a subsequence $(b_{n_{k_l}})_{l \in \mathbb{N}}$ of $(b_{n_k})_{k \in \mathbb{N}}$ which converges to $b_0 \in B$. The subsequence $((a_{n_{k_l}}, b_{n_{k_l}}))_{l \in \mathbb{N}}$ of $((a_n, b_n))_{n \in \mathbb{N}}$ obviously converges to $(a_0, b_0) \in A \times B$. \square

THEOREM 3.47. Any set of the form $[a_1, b_1] \times [a_2, b_2] \times \dots \times [a_n, b_n] \subset \mathbb{R}^n$ are compact.

Proof. Proof by induction using Lemma 3.46. \square

THEOREM 3.48. (HEINE–BOREL.) Consider the metric space \mathbb{R}^n equipped with one of the standard metrics d_1 , d_2 or d_∞ . Any $A \subset \mathbb{R}^n$ is compact if and only if A is closed and bounded.

Proof. If A is bounded it is contained in some set of the form $[a_1, b_1] \times [a_2, b_2] \times \dots \times [a_n, b_n] \subset \mathbb{R}^n$ which is compact by Theorem 3.47. Since A is therefore a closed subset of a compact set, we have A compact by Theorem 3.38. \square

REMARK 3.49. The continuous functions

$$f_n : [0, 1] \longrightarrow \mathbb{R}, \quad x \mapsto \begin{cases} 1, & \text{for } x \leq \frac{1}{n+1} \\ -n(n+1)x + n+1, & \text{for } \frac{1}{n+1} < x \leq \frac{1}{n} \\ 0, & \text{for } \frac{1}{n} < x \leq 1 \end{cases}$$

in $C([0, 1])$ have the properties $d(f_n, f_m) = 1$ if $n \neq m$ and $d(f_n, 0) = 1$. The set $A = \{f_n, \quad n \in \mathbb{N}\} \subset B_2(0)$ is bounded in $C([0, 1])$ and closed, since any convergent sequence in A converges to a limit in A (there are no convergent sequences in A). But A is not compact, since the open covering

$$\mathcal{U} = \{B_{\frac{1}{2}}(f_n)\}$$

contains no finite \mathcal{U} -subcovering of A .

As additional example let us consider \mathbb{R} with the discrete metric and $A = (0, 1)$, or \mathbb{R}^n with the metric $\tilde{d}_2 : (x, y) \mapsto \frac{d_2(x, y)}{1 + d_2(x, y)}$ and $A = \mathbb{R}^n$. In both cases A is bounded and closed but not compact.

THEOREM 3.50. A compact metric space (X, d) is complete.

THEOREM 3.51. Let (X, d_X) be compact, and $f : (X, d_X) \longrightarrow (Y, d_Y)$ be continuous. Then $\mathcal{R}_f = f(X)$ is compact in (Y, d_Y) .

To appreciate compactness some more, let us visit a *stronger* form of continuity.

DEFINITION 3.52. Let (X, d_X) and (Y, d_Y) be metric spaces. A function $f : X \longrightarrow Y$ is *uniformly continuous* on X , if for all $\varepsilon \in \mathbb{R} > 0$ exists $\delta > 0$ s.t. $d_Y(f(x), f(y)) < \varepsilon$ for all x, \tilde{x} with $d_X(x, \tilde{x}) < \delta$.

This is obviously equivalent to $\forall \varepsilon > 0 \exists \delta > 0$ s.t. $\forall x \in X \quad f(B_\delta(x)) \subseteq B_\varepsilon(f(x))$.

PROPOSITION 3.53. Any uniformly continuous function $f : (X, d_X) \longrightarrow (Y, d_Y)$ is continuous.

EXAMPLE 3.54.

- i. $f : \mathbb{R} \longrightarrow \mathbb{R}, \quad x \mapsto 2x$ is uniformly continuous.
- ii. $f : \mathbb{R}^+ \longrightarrow \mathbb{R}, \quad x \mapsto \frac{1}{x}$ is continuous but not uniformly continuous.

THEOREM 3.55. Any continuous function defined on compact metric spaces is uniformly continuous. That is, given a compact metric space (X, d_X) and continuous $f : (X, d_X) \longrightarrow (Y, d_Y)$, then f is uniformly continuous as well. (See homework problem 11.2.)

3.4. Connectedness

Again, we constrain ourselves to metric spaces.

DEFINITION 3.56. A metric space (X, d) is *connected* if X and \emptyset are the only *clopen*, that is, open and closed, subsets of X .

A *separation* of a metric space (X, d) is a pair of nonempty open subsets $U, V \subset X$ with $X = U \cup V$ and $\emptyset = U \cap V$.

Any subset A of the metric space (X, d) is connected if the metric space $(A, d|_{A \times A})$ is connected.

PROPOSITION 3.57. A metric space (X, d) is connected if and only if there exists no separation of X .

The most important result of this section is fairly elementary:

THEOREM 3.58. If (X, d) is connected and $f : (X, d_X) \longrightarrow (Y, d_Y)$ is continuous, then $\mathcal{R}_f = f(X)$ is connected.

REMARK 3.59. Using the fact that images of compacts under continuous transformations are compact and that images of connected sets under continuous transformations are connected, we can easily see that none of the sets

- i. $[0, 1] \subset \mathbb{R}$
- ii. $[0, 1) \subset \mathbb{R}$
- iii. $S^1 = \{z \in \mathbb{C} : |z| = 1\}$ in \mathbb{C}
- iv. The 8 set $S^1 \cup \{z \in \mathbb{C} : |z - 2i| = 1\}$ in \mathbb{C}

is homeomorphic to another set in the list.

THEOREM 3.60. Let us consider the real line \mathbb{R} with metric d_1 , d_2 , and d_∞ . The following are equivalent:

- i. The set $A \subset \mathbb{R}$ is connected.
- ii. For any $a, b \in A \subset \mathbb{R}$ and any $c \in \mathbb{R}$ with $a < c < b$ we have $c \in A$.
- iii. The set $A \subset \mathbb{R}$ is a (possibly unbounded) interval.

That is, connected sets in \mathbb{R} are exactly intervals and vice versa.

COROLLARY 3.61. Let $f : [a, b] \longrightarrow \mathbb{R}$ be continuous. Then exists $c, d \in \mathbb{R}$ with $f([a, b]) = [c, d]$.

THEOREM 3.62. (INTERMEDIATE VALUE THEOREM.) Let (X, d) be connected and $f : X \longrightarrow \mathbb{R}$ be continuous. Given any x_1, x_2 in X and $c \in \mathbb{R}$ with $f(x_1) < c < f(x_2)$, then exists $x \in X$ with $f(x) = c$.

THEOREM 3.63. Let $S_i, i \in I$ be a family of connected sets in a metric space (X, d) . If $\bigcap_{i \in I} S_i \neq \emptyset$, then $\bigcup_{i \in I} S_i$ is connected.

EXAMPLE 3.64. Open and closed balls in $(\mathbb{R}^n, d_i), i = 1, 2, \infty$ are connected. To see this, let A be an open or closed ball in (\mathbb{R}^n, d_{i_0}) , for some $i_0 \in \{1, 2, \infty\}$. For $x \in A$ consider

$$f_x : [0, 1] \longrightarrow \mathbb{R}^n, \quad t \mapsto tx + (1 - t)x_0.$$

The functions f_x are continuous and their ranges \mathcal{R}_{f_x} are therefore connected. The result follows from Theorem 3.63 since

$$A = \bigcup_{x \in A} \mathcal{R}_{f_x} \quad \text{and} \quad \bigcap_{x \in A} \mathcal{R}_{f_x} = \{x_0\} \neq \emptyset.$$

DEFINITION 3.65. A metric space (X, d) is called *totally disconnected* if for each $x \in X$ and $\epsilon > 0$ exists a clopen set A in X with $x \in A \subseteq B_\epsilon(x)$.

EXAMPLE 3.66. Cantor's middle third set is an uncountable set which is totally disconnected.

3.5. Sequences of functions, uniform convergence

In this section we shall discuss in detail the metric space $C(X)$ of continuous, complex valued functions defined on a compact metric space X .

The metric on $C(X)$ has been discussed in numerous homework problems.

DEFINITION 3.67. Let (X, d_X) be a metric space and let $B(X)$ be the set of all bounded, complex valued functions on X , that is,

$$B(X) = \{f : X \longrightarrow \mathbb{C} : \text{for } f \text{ exists } M \in \mathbb{R}^+ \text{ such that } |f(x)| \leq M \text{ for all } x \in X\}.$$

On $B(X)$ we can define the metric

$$d_\infty(f, g) = \sup\{|f(x) - g(x)| : x \in X\}.$$

The set of continuous, complex valued functions on X is denoted by $C(X)$. Note that (X, d) being compact implies that all continuous functions defined on X are bounded and we have $C(X) \subseteq B(X)$, and, therefore, $C(X)$ inherits the metric d_∞ from $B(X)$.

DEFINITION 3.68. Let (X, d_X) and (Y, d_Y) be metric spaces and let $f_n : X \longrightarrow Y$, $n \in \mathbb{N}$ be a sequence of functions mapping X to Y .

The sequence $(f_n)_{n \in \mathbb{N}}$ *converges pointwise* to $f_0 : X \longrightarrow Y$, if $\lim_{n \rightarrow \infty} f_n(x) = f_0(x)$ for all $x \in X$, that is, if $\lim_{n \rightarrow \infty} d_Y(f_n(x), f_0(x)) = 0$ for all $x \in X$.

The sequence $(f_n)_{n \in \mathbb{N}}$ *converges uniformly* to $f_0 : X \longrightarrow Y$, if for all $\epsilon > 0$ exists $N \in \mathbb{N}$ such that

$$d_Y(f_n(x), f_0(x)) < \epsilon \quad \text{for all } x \in X \text{ and for all } n \geq N.$$

That is

$$\lim_{n \rightarrow \infty} \sup \{d_Y(f_n(x), f_0(x)) : x \in X\} = 0.$$

PROPOSITION 3.69. The sequence (f_n) converges in $(B(X), d_\infty)$ to f_0 if and only if (f_n) converges to $f_0 : X \longrightarrow \mathbb{C}$ uniformly.

THEOREM 3.70. Let (f_n) be a sequence of continuous functions in $(B(X), d_\infty)$ which converges to f_0 . Then f_0 is continuous and for any sequence (x_k) in X with $\lim_{k \rightarrow \infty} x_k = x_0$ we have

$$\lim_{n \rightarrow \infty} \lim_{k \rightarrow \infty} f_n(x_k) = f_0(x_0) = \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} f_n(x_k).$$

COROLLARY 3.71. If (X, d) is compact, then $C(X)$ is a closed subspace of $B(X)$.

Proof. Since (X, d) is compact we have $f(X)$ is compact and therefore bounded for any continuous $f : X \longrightarrow \mathbb{C}$. Hence $C(X) \subseteq B(X)$ and, by Theorem 3.70 we have $C(X)$ closed in $(B(X), d_\infty)$. \square

THEOREM 3.72. Let (X, d) be a compact metric space. Then $(C(X), d_\infty)$ is a complete metric space.

4. DIFFERENTIATION

4.1. Central results

In this section, we shall discuss derivatives of real valued functions defined on subsets of \mathbb{R} . Our main objective is to illuminate the interplay of continuity and differentiability.

To define derivatives of real valued functions, we shall analyze so-called difference quotients. The discussion of such requires the following definition of functional limits.

DEFINITION 4.1. Let (X, d_X) and (Y, d_Y) be metric spaces and let f map X to Y . If x is a cluster point in X , we write $f(x) \rightarrow y_0$ as $x \rightarrow x_0$ or $\lim_{x \rightarrow x_0} f(x) = y_0$ if $y_0 \in Y$ and if for any $\epsilon > 0$ exists $\delta > 0$ such that $d_Y(f(x), y_0) < \epsilon$ whenever $0 < d_X(x, x_0) < \delta$. The point $y_0 \in Y$ is called functional limit of f as x approaches x_0 .

REMARK 4.2. If we restrict ourselves to cluster points, we could rephrase previous results using functional limits. E.g., we have:

- i. If x is a cluster point in (X, d_X) , then $\lim_{x \rightarrow x_0} f(x) = y_0$ if and only if for all sequences (x_n) in X with $x_n \neq x_0$, $n \in \mathbb{N}$, we have $\lim_{n \rightarrow \infty} f(x_n) = y_0$.
- ii. Let (X, d_X) and (Y, d_Y) be metric spaces, let f map X to Y , and let x be a cluster point in (X, d_X) . Then $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ if and only if f is continuous at x_0 .
- iii. For U open in \mathbb{R} we have $U' \supset U$, hence, the restriction to cluster points will not play a role in the following discussion of derivatives. By the way, any set A in a metric space (X, d) with $A = A'$ is called *perfect*.

DEFINITION 4.3. Let $A \subseteq \mathbb{R}$ and $f : A \rightarrow \mathbb{R}$. We say that f is *differentiable* at a cluster point x_0 in A , that is, at $x_0 \in A \cap A'$, if

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = L$$

for some $L \in \mathbb{R}$. In this case L is called derivative of f at x_0 and we write $f'(x_0) = L$. If $A \subseteq A'$ and f is differentiable at x for all $x \in A$, then we call f differentiable on A .

Further, we have that $f'(x_0) = L$ if and only if $\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} = L$.

In order to avoid “cluster point” disclaimers, we shall mostly restrict ourselves to consider open sets U as domains of differentiable functions. Open subsets of \mathbb{R} have the property that all its elements are cluster points.

EXAMPLE 4.4. For $\exp : \mathbb{R} \rightarrow \mathbb{R}$, $x \mapsto \sum_{n=0}^{\infty} \frac{x^n}{n!}$, we have $\exp'(x) = \exp(x)$.

Differentiable functions are continuous:

THEOREM 4.5. For U open in \mathbb{R} and $f : U \rightarrow \mathbb{R}$ differentiable at $x_0 \in U$ we have f continuous at x_0 .

THEOREM 4.6. (SUM, PRODUCT, AND QUOTIENT RULE.) Let U be open in \mathbb{R} and $f, g : U \rightarrow \mathbb{R}$ be differentiable at $x_0 \in U$. Then

- i. $f + g$ is differentiable at x_0 and $(f + g)'(x_0) = f'(x_0) + g'(x_0)$.
- ii. fg is differentiable at x_0 and $(fg)'(x_0) = f'(x_0)g(x_0) + f(x_0)g'(x_0)$.
- iii. If $g(x_0) \neq 0$, then $\frac{f}{g}$ is differentiable at x_0 and $\left(\frac{f}{g}\right)'(x_0) = \frac{f'(x_0)g(x_0) - f(x_0)g'(x_0)}{g(x_0)^2}$.

THEOREM 4.7. (CHAIN RULE.) Let U, V be open in \mathbb{R} and $f : U \rightarrow V$ be differentiable at $x_0 \in U$ and $g : V \rightarrow \mathbb{R}$ be differentiable at $f(x_0) \in V$. Then $g \circ f$ is differentiable at x_0 and we have $(g \circ f)'(x_0) = g'(f(x_0))f'(x_0)$.

EXAMPLES 4.8. For $n = 0, 1, 2, 3$ set $f_n(x) = \begin{cases} x^n \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$. Note that f_n , $n = 0, 1, 2, 3$, is continuous and differentiable on $\mathbb{R} \setminus \{0\}$, and its derivative is a continuous function on $\mathbb{R} \setminus \{0\}$.

- i. f_0 is not continuous at 0.
- ii. f_1 is continuous at 0 but not differentiable at 0.
- iii. f_2 is differentiable at 0, and, hence, on \mathbb{R} , but its derivative f_2' is not continuous at 0.
- iv. f_3 is again differentiable on \mathbb{R} and its derivative f_3' is continuous on \mathbb{R} .

THEOREM 4.9. INTERIOR EXTREMUM THEOREM. Let $U \subset \mathbb{R}$ be open and $f : U \rightarrow \mathbb{R}$ be differentiable on U . If there exists a maximum [resp. minimum] of f at c , then $f'(c) = 0$.

THEOREM 4.10. ROLLE'S THEOREM. Let $b > a$, and $f : [a, b] \rightarrow \mathbb{R}$ be continuous and differentiable on (a, b) . If $f(a) = f(b)$, then exists $c \in (a, b)$ such that $f'(c) = 0$.

THEOREM 4.11. MEAN VALUE THEOREM. Let $b > a$, and $f : [a, b] \rightarrow \mathbb{R}$ be continuous and differentiable on (a, b) . Then exists $c \in (a, b)$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.

THEOREM 4.12. GENERALIZED MEAN VALUE THEOREM. Let $b > a$, and $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous and differentiable on (a, b) . Then exists $c \in (a, b)$ such that $(g(b) - g(a))f'(c) = (f(b) - f(a))g'(c)$.

Proof. Apply Rolle's theorem to $h(x) = (g(b) - g(a))f(x) - (f(b) - f(a))g(x)$, $x \in [a, b]$. \square

We have seen that not all functions which are differentiable on an open interval have continuous derivatives. Nevertheless, they do not have "jump-discontinuities":

THEOREM 4.13. DARBOUX'S THEOREM. Let $f : (a, b) \rightarrow \mathbb{R}$ be differentiable. Then the function $f' : (a, b) \rightarrow \mathbb{R}$ has the intermediate value property, that is, for $u, v \in (a, b)$ and $\xi \in \mathbb{R}$ with $f'(u) < \xi < f'(v)$ exists $c \in (\min\{u, v\}, \max\{u, v\})$ with $f'(c) = \xi$.

DEFINITION 4.14. A function $f : A \longrightarrow \mathbb{R}$ is

- i. *monotonically increasing*, or simply *increasing*, if $f(x) \leq f(y)$ for all $x, y \in A$, with $x < y$
- ii. *strictly monotonically increasing*, or simply *strictly increasing*, if $f(x) < f(y)$ for all $x, y \in A$, with $x < y$
- iii. *monotonically decreasing*, or simply *decreasing*, if $f(x) \geq f(y)$ for all $x, y \in A$, with $x < y$, and
- iv. *strictly monotonically decreasing*, or simply *strictly decreasing*, if $f(x) > f(y)$ for all $x, y \in A$, with $x < y$.

A function is called *monotone* if it is either monotonically increasing or decreasing, and *strictly monotone* if it is either strictly increasing or strictly decreasing.

THEOREM 4.15. Let $f : (a, b) \longrightarrow \mathbb{R}$ be differentiable. Then f is

- i. monotonically increasing if and only if $f'(x) \geq 0$ for all $x \in (a, b)$, and
- ii. monotonically decreasing if and only if $f'(x) \leq 0$ for all $x \in (a, b)$.

EXAMPLE 4.16. Discussion of x^n , $n \in \mathbb{N}_0$, including the remark that $f(x) = x^3$ is strictly increasing on \mathbb{R} but $f'(0) = 0$.

THEOREM 4.17. Let $f : [a, b] \longrightarrow \mathbb{R}$ be continuous and strictly monotone. Let $[c, d] = f([a, b])$ and $\phi : [c, d] \rightarrow \mathbb{R}$ be the inverse function of f . If f is differentiable at $x_0 \in (a, b)$ with $f'(x_0) \neq 0$, then ϕ is differentiable at $y_0 = f(x_0) \in (c, d)$ and $\phi'(y_0) = \frac{1}{f'(x_0)} = \frac{1}{f'(\phi(y_0))}$.

DEFINITION 4.18. INFINITE LIMITS AND LIMITS AT INFINITY. Let $f : A \longrightarrow \mathbb{R}$, $A \subseteq \mathbb{R}$ and let $x_0, L \in \mathbb{R}^* = \mathbb{R} \cup \{+\infty, -\infty\}$. For $\epsilon > 0$, we call $(\frac{1}{\epsilon}, \infty)$ an ϵ -neighborhood of ∞ and $(-\infty, -\frac{1}{\epsilon})$ an ϵ -neighborhood of $-\infty$.

Further, we say that $f(x) \rightarrow L$ as $x \rightarrow a$ or $f(x)$ approaches L as x approaches x_0 , if for all $\epsilon > 0$ exists a $\delta > 0$ with

$$\left. \begin{array}{ll} x_0 \in A' \subset \mathbb{R} : & 0 < |x - x_0| < \delta \\ \text{or } x_0 = \infty : & x_0 > \frac{1}{\delta} \\ \text{or } x_0 = -\infty : & x_0 < -\frac{1}{\delta} \end{array} \right\} \text{ with } x \in A \text{ implies } \left\{ \begin{array}{ll} f(x) \in B_\epsilon(x_0), & \text{if } L \in \mathbb{R}; \\ f(x) \in (\frac{1}{\epsilon}, \infty) & \text{if } L = \infty; \\ f(x) \in (-\infty, -\frac{1}{\epsilon}), & \text{if } L = -\infty. \end{array} \right.$$

THEOREM 4.19. L'HOSPITAL'S RULE Suppose that f and g are real valued differentiable functions defined on (a, b) where $a \in \mathbb{R} \cup \{-\infty\}$ and $b \in \mathbb{R} \cup \{\infty\}$, $g'(x) \neq 0$ on (a, b) , and $\frac{f'(x)}{g'(x)} \rightarrow L \in \mathbb{R}^*$ as $x \rightarrow a$.

If $f(x) \rightarrow 0$ and $g(x) \rightarrow 0$ as $x \rightarrow a$, or if $g(x) \rightarrow \infty$ as $x \rightarrow a$, **then** $\frac{f(x)}{g(x)} \rightarrow L \in \mathbb{R}^*$ as $x \rightarrow a$.

An analogous statement holds of course if $x \rightarrow b$ or if $g(x) \rightarrow -\infty$.

4.2. Taylor series

DEFINITION 4.20. HIGHER DERIVATIVES. For $r \in \mathbb{N}$ we say that $f : U \rightarrow \mathbb{R}$, U open, has an n -th derivative at x_0 if $f^{(0)} = f$, $f^{(1)} = f'$, $f^{(2)} = f''$, \dots , $f^{(n-1)} = f^{(n-2)'}$ are defined on $(x_0 - \epsilon, x_0 + \epsilon)$ for some $\epsilon > 0$ and $f^{(n-1)}$ is differentiable at x_0 .

If f has an n -th derivative on U , that is, f has an n -th derivative at x_0 for all $x_0 \in U$, and if $f^{(n)} = f^{(n-1)'}$ is continuous on U , then we write $f \in C^n(U)$. If $f \in C^n(U)$ for all $n \in \mathbb{N}$, then we write $f \in C^\infty(U)$ and say f is called *smooth*.

Certainly, we shall also write $C^n(A)$ or $C^\infty(A)$ if A has the property that all its members are cluster points, that is, $A \subseteq A'$. For example, we could consider $C^2([0, 1])$.

REMARK 4.21. Note that the notation described above is in accordance to the symbol $C^0(U) = C(U)$ of continuous functions on U .

If U is an interval, for example $U = (a, b)$ we shall write $C^n(a, b)$ rather than $C^n((a, b))$.

THEOREM 4.22. TAYLOR'S THEOREM. Given $f : (a, b) \rightarrow \mathbb{R}$ and $n \in \mathbb{N}$ with $f \in C^{n-1}(a, b)$ and $f^{(n)}$ defined (but not necessarily continuous) on (a, b) . For x_0 in (a, b) define the $n - 1$ -th degree *Taylor polynomial* as

$$P_{f, x_0}(x) = \sum_{k=0}^{n-1} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k, \quad x \in (a, b).$$

For any $x \in (a, b)$ exists a ξ_x between x_0 and x such that

$$f(x) = P_{f, x_0}(x) + \frac{f^{(n)}(\xi_x)}{n!} (x - x_0)^n.$$

REMARK 4.23. Taylor's Theorem is used to compute approximate values of functions by means of evaluating polynomials.

For example, if $|f^{(n)}(\xi)| < M$ for all ξ between x and x_0 , then we have

$$|f(x) - P_{f, x_0}(x)| = \left| \frac{f^{(n)}(\xi_x)}{n!} (x - x_0)^n \right| \leq \frac{M}{n!} |x - x_0|^n$$

For x being close to x_0 the right hand side, and, therefore, the approximation error are small.

COROLLARY 4.24. If $f \in C^n(a, b)$ with $f^{(n)}(\xi) = 0$ for all $\xi \in (a, b)$, then f is a polynomial of degree at most $(n - 1)$.

COROLLARY 4.25. If for $f \in C^\infty(a, b)$ there exists $M > 0$ with $|f^{(n)}(\xi)| \leq M$ for all $\xi \in (a, b)$ and $n \in \mathbb{N}$, then for any $x_0 \in (a, b)$, we have

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k, \quad x \in (a, b).$$

DEFINITION 4.26. For $f \in C^\infty(a, b)$ and $x_0 \in (a, b)$, call the formal power series

$$T_{f,x_0}(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k, \quad x \in (a, b)$$

Taylor series of f at x_0 .

REMARK 4.27.

i. The radius of convergence of a Taylor series is not necessarily larger than 0.

ii. Even if the Taylor series of a function converges, it might not converge to the function.

For example, consider $f(x) = \begin{cases} e^{-\frac{1}{x^2}}, & \text{for } x \neq 0 \\ 0, & \text{else.} \end{cases}$ satisfies $f \in C^\infty(\mathbb{R})$, $f^{(n)}(0) = 0$ for

$n \in \mathbb{N}$ and, therefore, $T_{f,0}$ has radius of convergence $R = \infty$ and $T_{f,0}(x) = 0 \neq f(x)$ for $x \neq 0$.

THEOREM 4.28. Assume that (f_n) is a sequence of functions which are differentiable on (c, d) , and let $[a, b] \subset (c, d)$. If $\sum_{n=1}^{\infty} f_n(x)$ converges at some $x_0 \in [a, b]$ and $\sum_{n=1}^{\infty} f'_n(x)$ converges uniformly on $[a, b]$, then $\sum_{n=1}^{\infty} f_n(x)$ converges to a differentiable function, and

$$\left(\sum_{n=1}^{\infty} f_n(x) \right)' = \sum_{n=1}^{\infty} f'_n(x).$$

Proof. Use Theorem 3.70. □

PROPOSITION 4.29. If $f(x) = \sum_{k=0}^{\infty} c_k(x - x_0)^k$ for $x \in (a, b)$, then $f \in C^\infty(a, b)$ and $f^{(k)}(x) = c_k k!$ for $k \in \mathbb{N}$. Further, we have $f'(x) = \sum_{k=1}^{\infty} c_k k(x - x_0)^{k-1}$ for $x \in (a, b)$, that is, we can differentiate the series of functions f term by term.

Proof. Use Theorem 4.28. □

4.3. The exponential function and friends

The following theorem lists important facts regarding the exponential function $\exp(z) = \sum_{k=0}^{\infty} \frac{z^k}{k!}$, $z \in \mathbb{C}$, some of which we stated and proved earlier.

THEOREM 4.30. THE EXPONENTIAL FUNCTION.

- i. $\exp(z) = \sum_{k=0}^{\infty} \frac{z^k}{k!}$ converges absolutely for $z \in \mathbb{C}$.
- ii. $\exp(z + w) = \exp(z) \exp(w)$ for $z, w \in \mathbb{C}$.
- iii. $\exp(x) = \exp(1)^x = e^x$ for $x \in \mathbb{R}$.
- iv. $\exp'(x) = \exp(x)$ for $x \in \mathbb{R}$.
- v. $\exp(x) > 0$ for $x \in \mathbb{R}$ and \exp is strictly monotonically increasing.
- vi. $\exp(x) \rightarrow \infty$ as $x \rightarrow \infty$ and $\exp(x) \rightarrow 0$ as $x \rightarrow -\infty$.
- vii. $\exp : \mathbb{R} \rightarrow \mathbb{R}^+$ is bijective.
- viii. $\frac{x^n}{\exp(x)} \rightarrow 0$ as $x \rightarrow \infty$ for all $n \in \mathbb{N}$.

DEFINITION 4.31. The inverse function of $\exp : \mathbb{R} \rightarrow \mathbb{R}^+$ is called natural logarithm and is denoted by $\log : \mathbb{R}^+ \rightarrow \mathbb{R}$.

PROPOSITION 4.32.

- i. $\log(xy) = \log(x) + \log(y)$ for $x, y \in \mathbb{R}^+$.
- ii. The natural logarithm is a differentiable function with $\log'(x) = \frac{1}{x}$ for $x \in \mathbb{R}^+$.
- iii. For $x > 0$ we have $x^a = \exp(a \log(x)) = e^{a \log(x)}$ and $f : \mathbb{R}^+ \rightarrow \mathbb{R}$, $x \mapsto x^a$ is differentiable with $f'(x) = ax^{a-1}$.
- iv. For $a > 0$ we have again $a^x = \exp(x \log(a)) = e^{x \log(a)}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$, $x \mapsto a^x$ is differentiable with $g'(x) = a^x \log(a)$.

Proof. ii. Use Theorem 4.17, iii. and iv. by chain rule. □

DEFINITION 4.33. For $a > 0$, the function of $g(x) : \mathbb{R} \rightarrow \mathbb{R}^+$, $x \mapsto a^x$ is bijective and its inverse is called logarithm to base a . We shall denote g^{-1} by $\log_a : \mathbb{R}^+ \rightarrow \mathbb{R}$.

After discussing the behavior of the restriction of the function $\exp : \mathbb{C} \rightarrow \mathbb{C}$ to the real axis \mathbb{R} , that is, $\exp : \mathbb{R} \rightarrow \mathbb{C}$, we shall now consider its restriction to the imaginary axis $i\mathbb{R} \subset \mathbb{C}$. Once we described its properties, we fully understand $\exp : \mathbb{C} \rightarrow \mathbb{C}$ since $\exp(a + bi) = \exp(a) \exp(bi)$ for $a, b \in \mathbb{R}$.

We shall study $\exp : i\mathbb{R} \rightarrow \mathbb{C}$ by studying its real and imaginary part.

DEFINITION 4.34. We define the *sine* function $\sin : \mathbb{R} \longrightarrow \mathbb{R}$ by setting $\sin(x) = \operatorname{Im} \exp(ix)$ for $x \in \mathbb{R}$ and the *cosine* function $\cos : \mathbb{R} \longrightarrow \mathbb{R}$ by setting $\cos(x) = \operatorname{Re} \exp(ix)$ for $x \in \mathbb{R}$.

For convenience, we shall write $\cos x$ for $\cos(x)$, $\sin x$ for $\sin(x)$, $\cos^n x$ for $(\cos(x))^n$, and $\sin^n x$ for $(\sin(x))^n$, for $x \in \mathbb{R}$ and $n \in \mathbb{N}$.

THEOREM 4.35.

- i. $\sin x = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(2k+1)!}$ for $x \in \mathbb{R}$.
- ii. $\cos x = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k}}{(2k)!}$ for $x \in \mathbb{R}$.
- iii. $\sin' = \cos$ and $\cos' = -\sin$.
- iv. $\sin^2 x + \cos^2 x = 1$ for $x \in \mathbb{R}$.
- v. \cos and \sin are 2π -periodic, that is, $\sin(x + 2\pi) = \sin x$, $\cos(x + 2\pi) = \cos x$, where $\frac{\pi}{2}$ is the smallest $x > 0$ such that $\cos x = 0$.

COROLLARY 4.36. $\exp : \mathbb{C} \longrightarrow \mathbb{C}$ is $2\pi i$ -periodic.

Proof. $\exp(z + 2\pi i) = \exp(z) \exp(2\pi i) = \exp(z)(\cos(2\pi) + i \sin(2\pi)) = \exp(z)$ for $z \in \mathbb{C}$. \square

4.4. Fixed point theorems and approximative methods

DEFINITION 4.37. An element $x_0 \in X$ is called *fixed point* of $f : X \longrightarrow X$ if $f(x_0) = x_0$.

DEFINITION 4.38. A *contraction* on a metric space (X, d) is a map $f : X \longrightarrow X$ such that for some constant k , $0 \leq k < 1$, we have

$$d(f(x), f(y)) \leq k d(x, y) \text{ for all } x, y \in X.$$

PROPOSITION 4.39. Contractions are uniformly continuous mappings.

THEOREM 4.40. BANACH FIXED POINT THEOREM. If $f : X \longrightarrow X$ is a contraction on a complete metric space (X, d) , then exists a unique fixed point $x_0 \in X$, and for any choice of $x_1 \in X$, the sequence (x_n) defined by

$$x_1, x_2 = f(x_1), x_3 = f(x_2) = f(f(x_1)) = f \circ f(x_1), \dots, x_{n+1} = f(x_n), \dots,$$

converges to x_0 . Moreover, we have

$$d(x_n, x_0) \leq \frac{k}{1-k} d(x_k, x_{k-1}) \leq \frac{k^{n-1}}{1-k} d(x_2, x_1).$$

THEOREM 4.41. NEWTON'S METHOD.⁴ Let f be continuous on $[a, b]$ and twice differentiable on (a, b) . If $f(a) \leq 0$, $f(b) > 0$, $f'(x) \geq \delta > 0$ and $0 \leq f''(x) \leq M$ for $x \in (a, b)$, then exists a unique point $\xi \in (a, b)$ with $f(\xi) = 0$. Moreover, for any x_1 with $f(x_1) > 0$ the sequence recursively defined by

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

converges to ξ and we have

$$|x_{n+1} - \xi| \leq \frac{2\delta}{M} \left| \frac{M}{2\delta} (x_1 - \xi) \right|^{2n}.$$

THEOREM 4.42. Let $f : [a, b] \rightarrow \mathbb{R}$, $a < b$, be a differentiable function with $f([a, b]) \subset [a, b]$ and let $q < 1$ such that $|f'(x)| \leq q$, $\forall x \in D$. For $x_1 \in [a, b]$ set $x_n = f(x_{n-1})$ for $n \geq 1$. Then the sequence (x_n) converges to the unique solution $\xi \in D$ of the equation $f(\xi) = \xi$ and the following inequalities holds:

$$|\xi - x_n| \leq \frac{q}{1-q} |x_n - x_{n-1}| \leq \frac{q^n}{1-q} |x_1 - x_0|.$$

⁴We shall only give one of the many cases/versions of Newton's method.

- abelian group, 6
- absolute value of a complex number, 13
- absolutely convergent series, 21
- addition in a field, 7
- Algebraic Limit Theorem, 15
- antisymmetric relation, 4
- archimedean property, 8
- associativity, 6

- Banach Fixed Point Theorem, 45
- bijective, 3
- Bolzano–Weierstrass Theorem, 16
- boundary of a set, 29
- bounded above, 9
- bounded below, 9
- bounded sequences in metric spaces, 15
- bounded sets in metric spaces, 15

- Cantor set, 35
- Cartesian product, 3
- Cauchy Condensation Theorem, 22
- Cauchy criterion, 20
- Cauchy sequences, 19
- Cauchy–Schwarz Inequality, 25
- Chain Rule, 38
- clopen sets, 34
- closed ball, 25
- closed sets, 26, 29
- closure of a set, 29
- cluster points of a set, 29
- commutative group, 6
- compact set, 30
- complete metric spaces, 19
- complex numbers, 13
- conditionally convergent series, 21
- conjugate of a complex number, 13
- continuous functions on \mathbb{R} , 25
- continuous functions on metric spaces, 25
- continuous functions on topological spaces, 27
- contraction, 45
- convergence in the extended real number system, 17
- convergence in topological spaces, 28
- converging sequences in metric spaces, 14
- converging sequences in the reals, 14
- cosine function, 44
- countable sets, 12
- covering compact set, 30

Index

- covering of a set, 30

- Darboux’s Theorem., 38
- decreasing function, 39
- decreasing sequences, 15
- Dedekind–cut, 9
- differentiable functions, 37
- discrete metric, 14
- discrete topology, 27
- distance function, 14
- distributive law, 7
- domain of a function, 3
- domain of a relation, 3
- Dominated Convergence Theorem, 21

- e, 20
- equivalence class, 4
- equivalence relation, 4
- equivalent elements, 4
- euclidean metric, 14
- exponential function, 43
- extended real number system, 17

- field, 7
- finite cover, 30
- fixed point, 45
- function, 3
- functional limits, 37

- Generalized Mean Value Theorem, 38
- graph of a function, 3
- greatest lower bound, 9
- group, 6

- Heine–Borel Theorem, 32
- higher derivatives, 40
- homeomorph topological spaces, 28
- homeomorphism, 28

- identity element, 6
- imaginary part of a complex number, 13
- increasing function, 39
- increasing sequences, 15
- indiscrete topology, 27
- infimum, 9
- infinite limits, 17, 39
- infinite series, 20
- Inheritance Principle, 26
- injective, 3

- integers, 6
- Interior Extremum Theorem, 38
- interior of a set, 29
- intervals, 34
- inverse element, 6

- L'Hospital's Rule., 39
- least upper bound, 9
- least upper bound property, 10
- Lebesgue number, 31
- Leibniz Criterion for Alternating Series, 22
- limit inferior (liminf), 17
- limit point of a real valued sequence, 17
- limit superior (limsup), 17
- limits at infinity, 39
- linear Order, 7
- logarithm naturales, 43
- logarithm to base a , 43
- lower bound, 9

- mapping, 3
- Mean Value Theorem, 38
- metric, 14
- metric space, 14
- metric space of bounded functions, 36
- metric space of continuous functions, 36
- monotone function, 39
- monotone sequences, 15
- multiplication in a field, 7

- n -th derivative, 40
- n -th root, 12
- n -tupel, 3
- natural logarithm, 43
- neighborhood, 25
- neighborhood of $\pm\infty$, 39
- nested interval property, 12
- Newton's Method, 45

- one-to-one, 3
- onto, 3
- open ball, 25
- open cover, 30
- open sets, 26, 27
- Order Limit Theorem, 16
- order relation, 7
- ordered field, 8

- partial sums, 20
- partition of a set, 4

- pointwise convergence of sequences of functions, 36
- power series, 23
- product of series, 24
- Product Rule, 38

- Quotient Rule, 38

- radius of convergence, 23, 42
- range of a function, 3
- range of a relation, 3
- Ratio Test, 22
- rational numbers, 7
- real numbers, 9
- real part of a complex number, 13
- rearrangement of a series, 21
- reflexive relation, 4
- relation, 3
- representative of an equivalence class, 4
- Rolle's Theorem, 38
- Root Test, 22

- separation of a metric space, 34
- sequences, 12
- sequentially compact set, 30
- series, 20
- sine function, 44
- smooth functions, 40
- strictly decreasing function, 39
- strictly decreasing sequences, 15
- strictly increasing function, 39
- strictly increasing sequences, 15
- strictly monotone function, 39
- subcoverings, 30
- subsequence of a sequence, 16
- Sum Rule, 38
- supremum, 9
- surjective, 3
- symmetric relation, 4

- Taylor polynomial, 40
- Taylor series, 41
- Taylor's Theorem, 40
- topological space, 27
- topology on X , 27
- totally disconnected sets, 35
- transitive relation, 4
- trivial topology, 27

- uniform continuity, 33

uniform convergence of sequences of functions,
36
uniqueness of the real number system, 10
upper bound, 9