

$$\left\{ \begin{array}{c} \text{h}^{\frac{1}{2}} \nabla^+ \\ \text{h}^{-\frac{1}{2}} \nabla^+ \end{array} \right\} \xleftarrow[\Gamma]{} {}^{2^L} \mathbb{I}$$

$$\left\{ \begin{array}{c} \text{h}^{\frac{1}{2}} \nabla^+ \\ \text{h}^{-\frac{1}{2}} \nabla^+ \end{array} \right\} \ni \Gamma_B \text{ standard basis}$$

$$\mathbf{h}' = \Gamma \underbrace{\Gamma}_{\mathbf{h}}$$

$${}^A \delta_B = {}^A \Gamma \Gamma_B$$

$$\begin{array}{ccc} & & {}^{2^L} \mathbb{I} \\ & \searrow & \uparrow \\ \mathbf{h} \times & \left\{ \begin{array}{c} \text{h}^{\frac{1}{2}} \nabla^+ \\ \text{h}^{-\frac{1}{2}} \nabla^+ \end{array} \right\} & \mathbf{h}' \\ & \swarrow & \downarrow \\ & \mathbf{h}' = \mathbf{h} \underbrace{\mathbf{h}}_{\mathbf{h}'} & \mathbf{h}' \end{array}$$

$$\mathbf{h} \times \left\{ \begin{array}{c} \text{h}^{\frac{1}{2}} \nabla^+ \\ \text{h}^{-\frac{1}{2}} \nabla^+ \end{array} \right\} \ni \mathbf{h}_B \text{ basis}$$

$$\mathbf{h}' = \mathbf{h}' \underbrace{\mathbf{h}}_{\mathbf{h}'}$$

$${}^A \delta_B = {}^A \mathbf{h} \mathbf{h}_B$$