


$$\mathbb{R} \triangleleft \ni \mathbb{L} \supset \mathbb{K} \text{ rund}$$

$$\mathbb{K} \underset{\text{face}}{\subseteq} \mathbb{L} \in \text{HI}^0_0 \text{ stnM cpt}$$

$$\mathbb{K} \lrcorner \mathbb{K} \triangleleft_0 \mathbb{R} = \frac{\gamma \in \mathbb{K} \triangleleft_0 \mathbb{R}}{\mathbb{K} | \gamma = 0} \quad \text{o-ideal} \quad \mathbb{K} \triangleleft_0 \mathbb{R}$$



$$\mathbb{K} \underset{\mathbb{K}}{\subseteq} \mathbb{L} \Rightarrow \mathbb{K} \lrcorner \mathbb{K} \triangleleft_0 \mathbb{R} \sqsubset_{\mathbb{K}} \mathbb{K} \triangleleft_0 \mathbb{R}$$

$$\mathbb{K} \underset{\mathbb{K}}{\subseteq} \mathbb{L} \Rightarrow \mathbb{K} \lrcorner \mathbb{K} \triangleleft_0 \mathbb{R} \sqsubset_{\mathbb{K}} \mathbb{K} \triangleleft_0 \mathbb{R}$$